


Comparison of estradiol benzoate and cypionate in Girolando cows submitted to a timed artificial insemination

Comparação do benzoato e do cipionato de estradiol em vacas Girolando submetidas à inseminação artificial em tempo fixo

Otávia Reis e Silva¹ ; Marco Roberto Bourg de Mello²; Lara Nogueira Silencio¹; Joaquim Esquerdo Ferreira¹; Samuel Rodrigues Bonamichi do Couto¹; Rosane Scatamburlo Lizieire Fajardo³; Osvaldo de Almeida Resende⁴

¹ Universidade Federal Rural do Rio de Janeiro, Rio de Janeiro – RJ, Brazil

² Universidade Federal Rural do Rio de Janeiro, Instituto de Zootécnica, Departamento de Reprodução Animal, Rio de Janeiro – RJ, Brazil

³ Empresa de Pesquisa Agropecuária do Estado do Rio de Janeiro, Rio de Janeiro – RJ, Brazil

⁴ Empresa Brasileira de Pesquisa Agropecuária, Embrapa Agrobiologia, Rio de Janeiro – RJ, Brazil

ABSTRACT

The various studies on the manipulation of the estrous cycle culminated in the development of estrous and ovulation synchronization protocols, to the point of defining an optimum moment for artificial insemination. The induction of ovulation is a primordial part for the determination of the moment of the TAI, so to study which hormones and how they behave allows a better understanding of this physiological process and, to manipulate it efficiently. The aim of the present study was to compare the use of estradiol benzoate (EB) and cypionate (ECP) as ovulation inducer in Girolando cows submitted to Timed Artificial Insemination (TAI). Ovulations of 108 Girolando cows were synchronized and the initial day of treatment, recorded as D0, which was when the animals received an intravaginal device containing 1g of progestogen and an intramuscular application (im) of 2mg of EB. After 8 days (D8), the device was removed and 500µg of cloprostenol (im) was administered to all females. Then, the animals were randomly divided into two treatments: BG Group (n = 52) and CG Group (n = 56). The CG animals received 1mg of ECP (im) as the ovulation inducer at the time of device removal, while BG Group cows received 1mg of EB (im) 24 h later (D9). TAI occurred at D10. After device removal, ultrasound evaluations were performed every 12 h up to ovulation. The following reproductive parameters were evaluated: interval from intravaginal device removal to ovulation - IDO (hours); interval from ovulation to TAI-IOT (hours); diameter of the largest follicle at intravaginal device removal (mm); maximum diameter of the DF (mm) at D10; the dominant follicle growth rate (mm/day); synchronization rate - SR (%); ovulation rate - OR (%) and PR - pregnancy rate (%). Only the ovulation rate presented a statistical difference ($p < 0.05$). In conclusion, despite the ovulation rate difference, both estradiol esters administered were effective and presented similar pregnancy rates in Girolando cows submitted to TAI.

Keywords: Follicular dynamics. Estradiol esters. Induction of ovulation.

RESUMO

Os diversos estudos sobre manipulação do ciclo estral culminaram com o desenvolvimento de protocolos de sincronização do estro e da ovulação, a ponto de se definir um momento ótimo para a inseminação artificial. A indução da ovulação é parte primordial para a determinação do momento da IATF. Dessa forma, estudar quais hormônios e como eles se comportam permite compreender melhor esse processo fisiológico e, ainda, manipulá-lo de forma eficiente. O objetivo do presente estudo foi comparar o uso de benzoato de estradiol (BE) e cipionato (ECP) em vacas Girolando submetidas à inseminação artificial em tempo fixo (IATF). As ovulações de 108 vacas Girolando foram sincronizadas e no início do tratamento, D0, os animais receberam um dispositivo intravaginal contendo 1g de progestágeno e uma aplicação intramuscular (im) de 2mg de BE. No D8, o dispositivo foi removido e 500µg de cloprostenol (im) foram administrados a todas as fêmeas. Depois disso, os animais foram divididos aleatoriamente em dois tratamentos: Grupo GB (n = 52) e Grupo GC (n = 56). Os animais do Grupo GC receberam 1mg de ECP (im) como indutor de ovulação no momento da retirada do dispositivo, enquanto as vacas do Grupo GB receberam 1mg de BE (im) 24 horas depois (D9). IATF ocorreu em D10. Após a remoção do dispositivo, as avaliações ultrassonográficas foram realizadas a cada 12 horas até a ovulação. Os parâmetros reprodutivos avaliados foram: intervalo entre a retirada do dispositivo intravaginal de P4 e

a ovulação- IRD (horas); intervalo entre a ovulação e a IATF- IOI (horas); diâmetro do maior folículo na retirada do dispositivo intravaginal de P4 (mm); taxa de crescimento do FD-TCFD (mm/dia); taxa de sincronização - TS (%); taxa de ovulação- TO (%) e taxa de concepção - TC (%). Apenas a taxa de ovulação apresentou diferença estatística ($p < 0.05$). Em conclusão, apesar da diferença na taxa de ovulação, ambos os ésteres de estradiol administrados foram eficazes e apresentaram taxas de prenhez semelhantes em vacas Girolando submetidas a IATF.

Palavras-chave: Dinâmica folicular. Ésteres de estradiol. Indução da ovulação.

Correspondence to:

Otávia Reis e Silva
Universidade Federal Rural do Rio de Janeiro
Rodovia BR 465, Km 07, s/n, Zona Rural
CEP: 23890-000, Rio de Janeiro – RJ, Brazil
e-mail: otaviareis@hotmail.com

Received: February 21, 2019

Approved: June 03, 2019

How to cite: Silva OR, Mello MRB, Silenciato LN, Ferreira JE, Couto SRB, Fajardo RSL, Resende OA. Comparison of estradiol benzoate and cypionate in Girolando cows submitted to a timed artificial insemination. *Braz J Vet Res Anim Sci.* 2019;56(3):e155078. <https://doi.org/10.11606/issn.1678-4456.bjvras.2019.155078>

Introduction

The average Brazilian productivity in 2017 was 1963 liters/cow/year (Instituto Brasileiro de Geografia e Estatística, 2017), an inexpressive productivity. The challenge to increase productivity of the cows in Brazil is related to numerous factors such as breed, environment and nutrition. In short, the goal to obtain better and more productive cows will undoubtedly include the selection of genetically superior animals.

Artificial insemination is the main tool to promote genetic improvement in Brazilian herds, but the poor identification of the estrus results in low service and conception rates and, consequently, low reproductive efficiency.

The search for alternatives to circumvent these difficulties led to the development of timed artificial insemination (TAI). The need for protocols that synchronize ovulations in a practical, cheap and efficient way led researchers to study the different hormones available on the market. The association of hormones also reduces the number of days of handling the animals, which is a great advantage as it will reduce the impact not only on animal welfare, but also save on labor.

Among the hormones used, estradiol cypionate is already widely used in beef cattle herds, since its protocol synchronizes ovulation with less handling differently from estradiol benzoate. A decrease in management and

handling makes the process more practical, cheaper, less stressful to the animals and guarantees efficiency for the producer. There are several forms of estrogen, including 17- β -estradiol, biologically active molecule, estradiol benzoate and estradiol cypionate. Although all are estrogens, each produces distinct 17- β -estradiol profiles in the circulation due to peculiarities in the esterification of the molecule, thus leading to changes in absorption and, consequently, its metabolism (Souza et al., 2005). The administration of 17 β -estradiol causes a faster action, since it does not need to be metabolized and transformed in the organism, according to observations by Larson & Ball (1992).

Estradiol cypionate, which is formed by the esterification of propionic acid cyclopentane, results in low solubility in water and, therefore, slower release at the site of administration, prolonging the biological activity when compared to estradiol benzoate and 17 β estradiol (Vynckier et al., 1990). According to Pancarci et al. (2002), the administration of 2mg EC (intramuscular) promotes an increase in plasma estradiol concentrations 12 h after administration, peaks at 36 h after administration and remains elevated for more than 96 h. Therefore, studies that compare these different hormones seeking to simplify and reduce the costs of TAI protocols for dairy cows are very important. Such studies can ensure good or even better reproductive efficiency and encourage insemination in Brazil. Consequently, the production costs will reduce and the products will be more competitive and, at the same time, the reproductive efficiency of the national herd will improve. The hypothesis was that EC administration (at device removal) decreases cow handling without affecting conception rate of dairy cows submitted to TAI programs, compared to EB given 24 h after P4 device removal. The objective of this study was to compare the use of benzoate and estradiol cypionate in Girolando cows submitted to Timed Artificial Insemination.

Materials and Methods

The present experiment was approved by the Commission of Ethics in the Use of Animals of the Animal Science Institute of the Federal Rural University of Rio de Janeiro (CEUA/IZ/UFRRJ) under process nº. 23083.024213/2017-14.

The experiment was carried out in Seropédica, RJ. Geographically, the region is located at 22°45'53"S, 43°41'56"W

and is 33m above sea level. A total of 108 Girolando cows that were non-pregnant, multiparous, lactating, cyclic and acyclic with a mean body score of 3.1 ± 0.5 (range 1 to 5) and mean age 7.2 ± 2.2 years were used. All animals were previously submitted to a gynecological examination by rectal palpation with the aid of ultrasound for the detection of cyclicity and abnormalities of the reproductive tract before initiating hormonal protocols and monitoring follicular dynamics.

The hormonal protocols for TAI were initiated on a random day of the estrous cycle, and this day was recorded as day Zero (D0), when all cows received an intravaginal device with 1g of progestogen (DIB[®], Coopers, Sao Paulo, Brazil) and an intramuscular (im) application of 2.0 mg of estradiol benzoate (EB-ESTROGIN[®], Biofarm, Jaboticabal-SP, Brazil). After 8 days (D8), the intravaginal device was removed and 500µg of cloprostenol (CIOSIN[®], Intervet Schering Plow Animal Health, Sao Paulo, Brazil) was given intramuscularly (im) to all animals. At the same time, the animals were randomly divided into two experimental groups: Estradiol Cypionate Group (CG; n = 56) and Estradiol Benzoate Group (BG; n = 52). The cows in the CG received as the ovulation inducer 1.0 mg of estradiol cypionate intramuscular (ECP[®], Pfizer, Sao Paulo, Brazil) at the time of removal of the intravaginal device, and the cows in the BG received 1.0 mg of estradiol benzoate the morning of the following day (D9). TAI occurred on D10 and the CG cows were inseminated in the morning (48 h after DIB[®] removal) and BG in the afternoon (60 h after DIB[®] removal). All hormonal applications were done at the same time of day (08:00 AM).

Ultrasound examinations were performed using the transrectal technique with a 7.5MHz linear transducer (Mindray DP2200 vet) and started on D8 of the synchronization protocol, which is the same day the intravaginal device was removed. These evaluations were repeated every 12 h until 24 h after TAI. The moment of ovulation was defined as the time of disappearance of a previously identified dominant follicle (DF) from one ultrasound examination

to the next, minus 6 h. In each evaluation, the largest follicle with a diameter greater than 4 mm was measured in order to monitor follicular growth and determine maximum diameter of the dominant follicle (FD).

The following reproductive parameters were evaluated: interval from intravaginal device removal to ovulation - IDO (hours); interval from ovulation to TAI-IOT (hours); diameter of the largest follicle at intravaginal device removal (mm); maximum diameter of the DF (mm) at D10; the dominant follicle growth rate (mm/day); synchronization rate - SR (Synchronized animals/treated animals \times 100); ovulation rate - OR (ovulated animals/treated animals \times 100), and PR - pregnancy rate (pregnant animals/ treated animals \times 100). The diagnosis of gestation was performed by rectal palpation with the aid of the ultrasound device accompanied by a linear transrectal transducer of 7.5MHz (Mindray DP2200 vet), 30 days after insemination.

The experimental design was completely randomized. All data were tested for normality by the Anderson-Darling test with a significance level of 5%. For the description of the results, standard deviations and means (mean \pm standard deviation) were used. The differences between the groups were tested by the general linear model of variance (ANOVA) and the Kruskal-Wallis test. In the analysis of variance, the effects of the classification variables were analyzed using estradiol benzoate (EB) and estradiol cypionate (ECP). The chi-square test, with a significance level of 5%, was used to compare the differences that imply proportions, such as those found in synchronization, ovulation and pregnancy rates.

Results

The results found in the present study for follicular growth rate, dominant follicle diameters measured in D10, intervals from progesterone device removal to ovulation and intervals from ovulation to TAI for BG and CG cows are presented in Table 1. None of the parameters evaluated above presented statistical difference.

Table 1 – Follicular growth rate, dominant follicle (DF) diameter at D10, intervals from progesterone device removal to ovulation and from TAI to ovulation, after estradiol cypionate (CE) and benzoate (BE) treatments in Girolando cows

Treatments (CE×BE)	Follicular growth rate (mm/day)	DF diameter (mm)	Interval from ovulation to	
			Progesterone device removal (h)	TAI (h)
BG	1.26 ± 1.33^a (n = 49)	14.06 ± 2.26^a (n = 49)	67.91 ± 8.68^a (n = 47)	8.68 ± 6.94^a (n = 47)
CG	1.14 ± 1.27^a (n = 50)	13.11 ± 3.02^a (n = 50)	63.90 ± 10.49^a (n = 40)	16.05 ± 10.20^a (n = 40)

Values with different superscripts within the same column are different ($p \leq 0.05$). TAI = Timed Artificial Insemination; BG = benzoate; group; CG = cypionate group.

Table 2 presents the pregnancy rates after using estradiol benzoate and cypionate according to the different intervals from TAI and ovulation. There was only difference in conception rate in the 12- to 24 h-interval between TAI and ovulation ($p = 0.03$).

The mean follicle diameters at D10 for pregnant and non-pregnant cows of the BG and CG are presented in Table 3. No statistical difference was observed between these values.

Synchronization, ovulation and pregnancy rates according to EB or ECP in TAI protocols are presented in Table 4. Non-ovulating animals were not considered for the calculation of the interval from intravaginal progesterone device removal to ovulation and the interval from ovulation to TAI. The ovulation rate of benzoate group was higher than cypionate ($p = 0.01$). The synchronization and pregnancy rates showed no difference ($p > 0.05$).

Table 2 – Pregnancy rate according to interval from TAI to ovulation in Girolando cows submitted to TAI

Interval from TAI to Ovulation (h)	Pregnancy rate (%)		P Value
	BG (n = 46)	CG (n = 40)	
Ovulated	42.86 (3/7) ^a	100 (1/1) ^a	0.2
0-12 hours	65.38 (17/26) ^a	62.50 (10/16) ^a	0.8
12-24 hours	33.33 (4/12) ^a	91.67 (11/12) ^b	0.03
>24 hours	100 (1/1) ^a	36.36 (4/11) ^a	0.2

Values with different superscripts within the same row are different ($p \leq 0.05$). TAI = Timed Artificial Insemination; BG = benzoate group; CG = cypionate group.

Table 3 – Mean follicle diameter and standard deviation of the largest follicle evaluated at D10 as a function of the treatments with ECP and EB and the number of pregnant Girolando cows submitted to TAI

Treatments	Status	Mean follicle diameter and standard deviation
BG (n = 48)	Pregnant (n = 25)	14.17 ± 1.65 ^a
	Non-Pregnant (n = 23)	13.98 ± 2.84 ^a
CG (n = 52)	Pregnant (n = 26)	13.69 ± 2.23 ^a
	Non-Pregnant (n = 26)	12.41 ± 3.66 ^a

Values with different superscripts within the same column are different ($p \leq 0.05$). EB = estradiol benzoate; ECP = estradiol cypionate; TAI = Timed Artificial Insemination; BG = benzoate group; CG = cypionate group.

Table 4 – Synchronization, ovulation and pregnancy rates according to EB or ECP in TAI protocols

Reproductive parameters	BG (n = 52)	CG (n = 56)	p value
Synchronization rate (%)	98.08 (51/52) ^a	96.43 (54/56) ^a	0.6
Ovulation rate (%)	90.38 (47/52) ^a	71.43 (40/56) ^b	0.01
Pregnancy rate (%)	48.08 (25/52) ^a	46.43 (26/56) ^a	0.8

Values with different superscripts within the same row are different ($p \leq 0.05$). EB = estradiol benzoate; ECP = estradiol cypionate; TAI = Timed Artificial Insemination; BG = benzoate group; CG = cypionate group; p value = probability of obtaining a test statistic equal to or more extreme than that observed in a sample.

Discussion

To our knowledge, this is the first report studying the effects of distinct ovulation inducers in estradiol and progesterone-based protocols on Girolando females submitted to TAI.

Studies from Borges et al. (2004), using Gir cows, demonstrated follicular growth rates ranging from 1.2 to 1.4 mm/day in animals that were not submitted to TAI protocols, which means that the results obtained in the present study approximate the values of a normal estrous cycle in Gir cows. On the other hand, Ferraz et al. (2013) obtained the highest follicular growth rate when they used GnRH as an ovulation inducer, obtaining a value of 1.07 ± 0.3 mm/day in Nelore cows, and the group that used estradiol benzoate as an inducer had a follicular growth rate of 0.58 ± 0.3 mm/day. Tortorella et al. (2017), working with *Bos taurus* females (cows and heifers), found a rate of 0.49 ± 0.02 mm/8 h for cows and 0.47 ± 0.06 mm/8 h for heifers. The dominant follicle growth rate appears to be higher in *Bos taurus* cows (Ginther, 2018) than in *Bos indicus* cows (Coutinho et al., 2007). The values found in the present study are intermediate to those found in the literature and can be explained since the cows used are Girolando (crossbred from *Bos taurus* x *Bos indicus*).

The variation in the values of follicular growth rates can be influenced by the subspecies to which the animals belong (*Bos taurus* x *Bos indicus*) or animal category (cows x heifers) (Hadiya et al., 2016). The use of different ovulation inducers did not appear to be a relevant factor in follicular growth rate. Plasma concentrations of progesterone seem to have a more significant influence than the ovulation inducer used, as demonstrated by Savio et al. (1993) who, by monitoring the estrous cycle of cows, demonstrated that high levels of progesterone decrease the follicular growth rate and, consequently, the size of the dominant follicle.

There was no significant difference between the diameter of the dominant follicle at D10 for the BG and CG groups. Andrade et al. (2012), using estradiol esters to induce ovulation in Nelore cattle, observed that the follicle diameter did not differ between the treatments that they used (ECP D8 x EB D9), obtaining, respectively, 13.03 ± 2.24 mm and 12.40 ± 1.34 mm. This result is like Sales et al. (2012), who found equivalent follicular diameters for treatment with estradiol cypionate (13.90 ± 0.40 mm) and estradiol benzoate (13.10 ± 0.40 mm) for the Nelore breed.

The results presented in the present study resemble other studies that have worked with dairy cows such as Franca et al. (2015), who used crossbred Holstein x Gir cows, and showed no difference related to the mean follicular

diameter for animals receiving ECP at D8 (10.71 ± 2.43 mm) and EB at D9 (11.45 ± 2.34 mm). More recently, crossbred cows showed 9.77 ± 1.36 mm and 10.97 ± 1.24 mm (Ramos Sobreira et al., 2017) for EB and ECP, respectively, with no statistical difference. Although the results obtained in the present study demonstrate a numerically superior means to the other studies mentioned, the similarity between the follicular diameters obtained with the use of both ovulation inducers suggests that the two were effective to stimulate the growth and also to obtain follicles greater than 10 mm which, according to Sartori et al. (2001), is a size that already has the capacity to respond to the action of LH and, consequently, to ovulate.

In the present study, EB produced ovulations at a similar time interval compared to ECP as demonstrated by Sales et al. (2012) and Callejas et al. (2016). In 2016, some researchers, who were evaluating the plasma concentrations of 17β -estradiol, reported that the application of ECP at the same time as the device removal and the application of EB one day after the withdrawal of the device generate similar hormonal profile. This characteristic occurs due to the pharmacokinetics of estradiol esters. Estradiol benzoate has a shorter half-life, and it induces an increase in the biologically active molecule (17β -estradiol) more rapidly, which explains why BE is administered 1 day after device removal and not at the same time.

Regarding the interval from TAI to ovulation, it was possible to observe that there was a statistical difference only between the BG and CG groups in the interval of 12-24 h, that is, the animals of the CG group that ovulated in this interval had a higher pregnancy rate when compared to BG.

Dransfield et al. (1998) and Roelofs et al. (2005) reported in their studies that the chance of an animal becoming pregnant increases as AI approaches the time of ovulation. Thus, in order to maximize the contact of the sperm with the oocyte, the AI should be performed close to the time of ovulation so that the egg does not age waiting for the spermatozoa. Based on the studies cited above and from other authors, the best time for AI would probably be between 16 and 26 h before ovulation (Dransfield et al., 1998; Roelofs et al., 2005). Therefore, the highest pregnancy rate in the 12- to 24 h-interval between AI and ovulation for CG is due to the fact that cows ovulate at a more favorable conception period. For example, in CG, 57.5% of the cows ovulated after 12 h in relation to AI, different from BG, where 71.73% of cows ovulated in the first 12 h after AI. This ovulation closest to AI is probably associated with the half-life of estradiol benzoate as explained above.

Success in TAI programs can be influenced by several factors, including the diameter of the dominant follicle at the time of artificial insemination (Pfeifer et al., 2015). The follicle diameter at the time of TAI is an important success factor in pregnancy rates, since larger follicles contain higher concentrations of estradiol and, therefore, a greater possibility of ovulation; consequently, there is an increase in the fertility indexes in synchronized herds (Sá et al., 2010). In addition, larger follicles give rise to a larger corpus luteum, thereby increasing the chances of establishing and maintaining pregnancy (Lonergan et al., 2013). Thus, a high concentration of pre-ovulatory estradiol generated by the greater diameter of the ovulatory follicle may influence the fertilization of females by promoting changes in the uterine environment, improving sperm transport and favoring conception (Sá et al., 2010).

Several researchers have reported the importance of follicular diameter on the day of insemination (Fujita et al., 2013; Ribeiro et al., 2013; Silveira et al., 2016). Silveira et al. (2016), working with the Nelore breed, found larger follicular diameters in the cows that conceived (13.26 mm \times 10.99 mm). Pfeifer et al. (2015) proposed to evaluate the size of the follicle on the day of TAI as a way to improve the fertility of the cows that will be inseminated and, depending on the size of the dominant follicle, it might be necessary to wait a few more hours and delay the TAI in order to optimize the pregnancy rate of the herd.

The present study did not observe a statistical difference between the follicular diameter on the day of the TAI and the pregnancy rate, since we did not use any other additional hormone with the objective of optimizing follicular growth, which is different from most of the above mentioned studies. Most of the researchers used eCG, a hormone that acts directly on the final follicle growth, generating a follicle with a larger diameter (Baruselli et al., 2008; Maciel et al., 2017).

Only the ovulation rate presented a difference between the groups. Estradiol benzoate was better than estradiol cypionate to induce ovulation. This may be associated with the fact that EB promotes higher concentrations of 17β -estradiol in the bloodstream earlier than the ECP, thus contributing to better ovulation rates, as discussed by Souza et al. (2005).

Despite the difference in ovulation rates, no statistical significance was observed in the pregnancy rates. This fact is possibly related to the CG cows, where these animals ovulated in a more propitious period for conception. So, despite the fact that the group presented lower ovulation rates, this was not sufficient to affect the pregnancy rates, since ovulations occurred at a more appropriate time than the BE group animals.

Conclusion

Although estradiol benzoate is more efficient in inducing ovulation than estradiol cypionate, both can be used with the same efficiency in TAI programs for Girolando cows because they produce the same pregnancy rates.

Conflict of Interest

The authors have no competing interest.

Ethics Statement

The present investigation attended all of the rules proposed by Brazilian National Council of Animal Experimentation (CONCEA) for controlling the use of animals for teaching

and research and was also approved by the Ethic Commission of Animals Usage in Experiments (CEUA) from the Federal University of Rio de Janeiro, Brazil.

Acknowledgements

The authors would like to thank the Federal Rural University of Rio de Janeiro, and the Agrolivestock Research Company of the State of Rio de Janeiro (PESAGRO). This work was conducted during a scholarship supported and financed by CAPES – Brazilian Federal Agency for Support and Evaluation of Graduate Education within the Ministry of Education of Brazil.

References

- Andrade BHDA, Ferraz PA, Rodrigues AS, Loiola MVG, Chalhoub M, Ribeiro AL Fo. Eficiência do cipionato de estradiol e do benzoato de estradiol em protocolos de indução da ovulação sobre a dinâmica ovariana e taxa de concepção de fêmeas nelore inseminadas em diferentes momentos. *Arch Vet Sci.* 2012;17(4):12. <http://dx.doi.org/10.5380/avs.v17i4.25710>.
- Baruselli PS, Jacomini JO, Sales JNS, Crepaldi GA. Importância do emprego da eCG em protocolos de sincronização para IA, TE e SOV em tempo fixo. In: *Anais do 3º Simpósio Internacional de Reprodução Animal Aplicada*; 2008; Londrina, PR. Londrina: SIRAA; 2008. p. 146-67.
- Borges AM, Torres CAA, Rocha VR Jr, Ruas JRM, Gioso MM, Fonseca JE, Carvalho GR, Maffili VV. Dinâmica folicular e momento da ovulação em vacas não lactantes das raças Gir e Nelore durante duas estações do ano. *Arq Bras Med Vet Zootec.* 2004;56(3):346-54. <http://dx.doi.org/10.1590/S0102-09352004000300010>.
- Callejas S, Uslenghi G, Larghi J, Cabodevila J. Comparación de dos protocolos para sincronizar ovulación e implementar inseminación artificial en vaquillonas. *Rev Vet.* 2016;25(2):100-4.
- Coutinho GTRM, Viana JHM, Sá WF, Camargo LS, Ferreira AM, Palhão PM, Nogueira LAG. Avaliação ultrasonográfica da dinâmica folicular e lútea em vacas da raça Guzerá Ultrasonographic evaluation of follicular dynamics and luteal function in Guzerá cows. *Arq Bras Med Vet Zootec.* 2007;59(5):1089-96. <http://dx.doi.org/10.1590/S0102-09352007000500001>.
- Dransfield MBG, Nebel RL, Pearson RE, Warnick LD. Timing of insemination for dairy cows identified in estrus by a radiotelemetric estrus detection system. *J Dairy Sci.* 1998;81(7):1874-82. [http://dx.doi.org/10.3168/jds.S0022-0302\(98\)75758-3](http://dx.doi.org/10.3168/jds.S0022-0302(98)75758-3). PMID:9710754.
- Ferraz PA, Rodrigues AS, Loiola MG, Lima MCC, Bittencourt TCBS, Ribeiro ADL Fo. Dinâmica folicular e taxa de concepção de vacas nelore sincronizadas com diferentes protocolos de indução da ovulação. *Arch Vet Sci.* 2013;18(4). <http://dx.doi.org/10.5380/avs.v18i4.31413>.
- Franca LM, Rodrigues AS, Brandão LGN, Loiola MVG, Chalhoub M, Ferraz PA, Bittencourt RF, Jesus EO, Ribeiro AL Fo. Comparison of two esters estradiol as inductors on ovulation follicular diameter and pregnancy rate of dairy cattle subject to program of FTAI. *Rev Bras Saúde Prod Anim.* 2015;16(4):958-65. <http://dx.doi.org/10.1590/S1519-99402015000400019>.
- Fujita AS, Weiss RR, Rossi P Jr, Kozicki LE, Greselle FVN, Bertol MAF. Taxa de gestação em novilhas nelore sincronizadas para IATF e inseminadas com sêmen resfriado e congelado. *Arch Vet Sci.* 2013;18(3). <http://dx.doi.org/10.5380/avs.v18i3.27935>.
- Ginther OJ. Variations in follicle-diameter deviation and a growth spurt in the dominant follicle at deviation in *Bos taurus* heifers. *Anim Reprod Sci.* 2018;188:155-64. <http://dx.doi.org/10.1016/j.anireprosci.2017.11.020>. PMID:29224966.
- Hadiya KK, Dhama AJ, Patel JA, Lunagariya PM, Patel AC. Follicular dynamics during estrous cycle in postpubertal and postpartum gir cattle. *Indian J Anim Reprod.* 2016;37(1):1-4.
- IBGE: Instituto Brasileiro de Geografia e Estatística. *Produção pecuária municipal*. Rio de Janeiro: IBGE; 2017. v. 45, p. 1-8.

- Larson LL, Ball P. Regulation of estrous cycles in dairy cattle: a review. *Theriogenology*. 1992;38(2):255-67. [http://dx.doi.org/10.1016/0093-691X\(92\)90234-I](http://dx.doi.org/10.1016/0093-691X(92)90234-I). PMID:16727134.
- Lonergan P, O'Hara L, Forde N. Role of diestrus progesterone on endometrial function and conceptus development in cattle. *Anim Reprod*. 2013;10(3):223-7.
- Maciel EB, Santos LHF, Santos KJG, Santos APP, Lopes JCS, Paula RS, Ferreira PCS, Pereira CC, Nunes CF Jr, Ramos PHS. Importância da utilização de eCG em protocolos de TAI em bovinos. *Anais da Semana do Curso de Zootecnia (SEZUS)*. 2017;10(1).
- Pancarci S, Jordan M, Risco ER, Schouten CA, Lopes MJ, Moreira FL, Thatcher W. Use of estradiol cypionate in a presynchronized timed artificial insemination program for lactating dairy cattle. *J Dairy Sci*. 2002;85(1):122-31. [http://dx.doi.org/10.3168/jds.S0022-0302\(02\)74060-5](http://dx.doi.org/10.3168/jds.S0022-0302(02)74060-5). PMID:11860104.
- Pfeifer LFM, Castro NA, Melo VTO, Neves PMA, Cestaro JP, Schneider A. Timed artificial insemination in blocks: a new alternative to improve fertility in lactating beef cows. *Anim Reprod Sci*. 2015;163:89-96. <http://dx.doi.org/10.1016/j.anireprosci.2015.10.002>. PMID:26481047.
- Ramos Sobreira R, Almeida IC, Oliveira FA, Siqueira JB, Barioni G, Lima DV, Siqueira LA. Cipionato de estradiol e benzoato de estradiol em protocolos de inseminação artificial em tempo fixo em novilhas mestiças. *Vet Zootec*. 2017;24(3):581-91. <http://dx.doi.org/10.35172/rvz.2017.v24.295>.
- Ribeiro ADL Fo, Ferraz PA, Rodrigues AS, Bittencourt TCBDS, Loiola MVG, Chalhoub M. Diâmetro do folículo no momento da inseminação artificial em tempo fixo e taxa de concepção em vacas Nelore. *Cienc Anim Bras*. 2013;14(4):501-7. <http://dx.doi.org/10.5216/cab.v14i4.19678>.
- Roelofs JB, Van Eerdenburg FJCM, Soede NM, Kemp B. Various behavioral signs of estrous and their relationship with time of ovulation in dairy cattle. *Theriogenology*. 2005;63(5):1366-77. <http://dx.doi.org/10.1016/j.theriogenology.2004.07.009>. PMID:15725444.
- Sá MFD Fo, Crespilho AM, Santos JEP, Perry GA, Baruselli PS. Ovarian follicle diameter at timed insemination and estrous response influence likelihood of ovulation and pregnancy after estrous synchronization with progesterone or progestin-based protocols in suckled *Bos indicus* cows. *Anim Reprod Sci*. 2010;120(1-4):23-30. <http://dx.doi.org/10.1016/j.anireprosci.2010.03.007>. PMID:20395079.
- Sales JNS, Carvalho JBP, Crepaldi GA, Cipriano RS, Jacomini JO, Maio JRG, Souza JC, Nogueira GP, Baruselli PS. Effects of two estradiol esters (benzoate and cypionate) on the induction of synchronized ovulations in *Bos indicus* cows submitted to a timed artificial insemination protocol. *Theriogenology*. 2012;78(3):510-6. <http://dx.doi.org/10.1016/j.theriogenology.2012.02.031>. PMID:22503845.
- Sartori R, Fricke PM, Ferreira JC, Ginther OJ, Wiltbank MC. Follicular deviation and acquisition of ovulatory capacity in bovine follicles. *Biol Reprod*. 2001;65(5):1403-9. <http://dx.doi.org/10.1095/biolreprod65.5.1403>. PMID:11673256.
- Savio JD, Thatcher WW, Badinga L, De La Sota RL, Wolfenson D. Regulation of dominant follicle turnover during the oestrous cycle in cows. *Reproduction*. 1993;97(1):197-203. <http://dx.doi.org/10.1530/jrf.0.0970197>. PMID:8464011.
- Silveira RO, Santos GM, Silveira CO, Maitan PP. Avaliação do tamanho do folículo ovulatório e da taxa de concepção de vacas Nelore em protocolos de IATF. *ANAIS SIMPAC*. 2016;6(1).
- Souza AH, Cunha AP, Caraviello DZ, Wiltbank MC. Profiles of circulating estradiol-17 β after different estrogen treatments in lactating dairy cows. *Anim Reprod*. 2005;2(4):224-32.
- Tortorella RD, Nogueira GP, Modesto MR, Silva PCP, Silva JP, Neves JP, Ramos AF. Characterizing emergence and divergence in the first follicular wave in a tropically adapted *Bos taurus* breed. *Theriogenology*. 2017;88:9-17. <http://dx.doi.org/10.1016/j.theriogenology.2016.09.041>. PMID:27865418.
- Vynckier L, Debackere M, Kruif A, Coryn M. Plasma estradiol-17 β concentrations in the cow during induced estrus and after injection of estradiol17 β benzoate and estradiol-17 β cypionate-a preliminary study. *J Vet Pharmacol Ther*. 1990;13(1):36-42. <http://dx.doi.org/10.1111/j.1365-2885.1990.tb00745.x>. PMID:2319634.

Financial Support: None.

Authors Contributions: Otávia Reis e Silva, Marco Roberto Bourg de Mello, Lara Nogueira Silenciato, Joaquim Esquerdo Ferreira, Samuel Rodrigues Bonamichi do Couto, Rosane Scatamburlo Lizieire Fajardo and Osvaldo de Almeida Resende participated in the writing of the article, considering the reading and review stages.