

Prevalence of postural deviations in the spine in schoolchildren: a systematic review with meta-analysis

Ana Paula Kasten¹, Bruna Nichele da Rosa¹, Emanuelle Francine Detogni Schmit¹, Matias Noll¹, Cláudia Tarragô Candotti¹

Doi: <http://dx.doi.org/10.7322/jhgd.127684>

Abstract

Introduction: The Information Age has had a major impact on citizens' lives. The invention of machinery, automobile, television and computer have induced individuals to adopt the "sitting" body posture in order to adapt to new technological needs.

Objective: To estimate the prevalence of spine postural deviations in Brazilian schoolchildren.

Methods: Searches were conducted in databases EMBASE, LILACS, PubMed, SCOPUS, SciELO, Science Direct, and Web of Science, as well as manual searches to identify studies that evaluated the prevalence of spine postural deviations in Brazilian schoolchildren. Two independent reviewers realized the study selection, evaluated the methodological quality and the risk of bias and extracted data. The homogeneity between the studies was evaluated and the quality of evidence level using the GRADE system.

Results: 29 studies were included, of which extracted the frequency of positive events to changes in cervical, thoracic and lumbar spine, as well as the frequency of scoliosis between schoolchildren. Even performing the meta-analysis separated by subgroups according to the spine region, the heterogeneity level it was up to 90%, it is not possible to perform the meta-analysis to estimate the prevalence of spine postural deviations in Brazilian schoolchildren.

Conclusion: There is low strength of evidence to establish a consensus about the values of the prevalence of spine postural deviations in Brazilian schoolchildren.

Keywords: posture, spine, child, adolescent.

INTRODUCTION

The Information Age has had a major impact on citizens' lives¹. The invention of machinery, automobile, television and computer have induced individuals to adopt the "sitting" body posture in order to adapt to new technological needs. Also, the increasing rate of adhesion of individuals to the new demands of convenience comfort has been an important causal factor of poor posture and back pain. Thus, both inadequate and sedentary habits, since childhood, contribute to the appearance of muscle weakness and ligament laxity, overloading the spine, which results in suffering, pain and disability²⁻⁵.

In addition, it has been accepted that postural problems related to changes in body shape often have their origin in childhood, especially those related to the spine⁶. In this sense, the identification or diagnosis of postural de-

viations during childhood and adolescence is of great importance in this phase of body growth and development, since the correction of these changes is mainly based on global posture reeducation. In fact, investing in reeducation during childhood and adolescence tends to minimize the need for a future conservative treatment that only aims to improve symptoms, since, after adolescence, bone growth has already ceased^{7,8}.

Some factors are preponderant in the causes of postural deviations in school-age children, such as, hours sitting in inappropriate positions in the classroom, inadequate seats, carrying school supplies in an inadequate way⁹⁻¹¹ and / or with weight above 10% of body weight⁵⁻¹⁰. In addition, the increase in Body Mass Index (BMI) in students diagnosed as pre-obese and obese results in a higher prevalence of asymmetry in the anterior, posterior and lateral planes¹²⁻¹⁴. In fact, these factors are evident world-

¹ Federal University of Rio Grande do Sul.

Corresponding author: Ana Paula Kasten - E-mail: anapaulakasten@hotmail.com

Suggested citation: Kasten AP, Rosa BN, Schmit EFD, Noll M, Candotti CT. Prevalence of postural changes in the spine in schoolchildren: a systematic review with meta-analysis. *J Hum Growth Dev.* 2017; 27(1): 99-108. Doi: <http://dx.doi.org/10.7322/jhgd.127684>

Manuscript submitted 2016, accepted for publication Jun 2016.

wide, in both developed and developing countries. In Iran, a developing country, two studies found a higher prevalence of changes in the spine of girls associated with sedentary lifestyle and excess weight carried in backpacks during the carrying of school supplies^{15,16}. In Japan, a developed country, a longitudinal study found an inverse relationship between paravertebral muscle strength and lumbosacral postural change. That is, during a 10-year follow-up, there was both a decrease in paravertebral muscle strength and an increase of around 10° in the lumbosacral angle, suggesting the need to promote activities to maintain muscle strength from childhood and adolescence^{17,18}.

Regarding developing countries, although Brazil stands out for the amount of research on the subject of school posture, as far as we know, no study was conducted synthesizing the Brazilian reality about the prevalence of postural deviations, especially those related to the spine. Thus, considering that knowledge of the profile of postural deviations in schoolchildren and adolescents is essential to assist in the development of public policies and intervention strategies, the objective of this study is to estimate the prevalence of postural deviations in Brazilian schoolchildren's spine through a systematic review with meta-analysis.

Table 1: Search strategy - PubMed

#1	("Students"[Mesh] OR "Students" OR "schoolchildren")
#2	("Spinal Curvatures"[Mesh] OR "Spinal Curvatures" OR "Hyperkyphosis"[Mesh] OR "Hyperkyphosis" OR "Lordosis"[Mesh] OR "Lordosis" OR "Scoliosis"[Mesh] OR "Scoliosis")
#3	("Prevalence"[Mesh] OR "Prevalence")
#4	(#1 AND #2 AND #3)

Eligibility Criteria

Two independent evaluators selected the potentially relevant studies from the titles, abstracts and full text. The studies selected for reading in full were evaluated according to the following eligibility criteria: (a) observational study; (B) exclusive sample of Brazilian schoolchildren aged 6 to 18 years; (C) to address the prevalence of postural deviations in the spine. Discordant cases were resolved by consensus²³.

Assessment of Studies

The evaluators obtained the study data independently and in a standardized form according to the following information: authors, year of publication, place of the holding of the study, sample evaluated, methods used and prevalence of changes in the spine.

Likewise, the studies were evaluated with respect to methodological quality and risk of bias through the Prevalence Critical Appraisal Instrument²⁴, an instrument developed to evaluate the methodological quality of studies that present prevalence data. This instrument consists of 10 items, which must be filled in as Yes, No, Unclear or Not Applicable, and then the items filled with Yes are punctuated. In this perspective, the evaluation of each study can present a score between 0 and 10, and the higher the score, the better the methodological evaluation of the study. In order to graduate and establish

METHODS

Type of study

This study comprised a systematic review of the literature¹⁹, registered in PROSPERO under the code CRD42015026504, and guided according to the recommendations of the Joanna Briggs Institute Reviewers Manual (The Systematic Reviewer of Prevalence and Incidence Data)²⁰, the guidelines of Collaboration Cochrane²¹ and the MOOSE (Meta-Analysis of Observational Studies in Epidemiology) report²².

Search strategy

In order to achieve the proposed objective, we conducted searches from September 23 to 25, 2015, in the EMBASE, LILACS, PubMed, SCOPUS, SciELO, Science Direct, and Web of Science databases. The terms and Boolean operators used were: "students" [AND] "spinal curvatures" [AND] "prevalence". No restriction of language and year of publication was done in the search, which was performed from the beginning of the bases until the moment of the search. The studies should be of the observational type. Manual searches were also conducted in the references of included studies. Table 1 presents the search strategy used in the PubMed database, and in the other databases this strategy was adapted according to the guidelines of each database.

categorical divisions of quality, a cutoff point was stipulated, with the minimum score of seven as the divisor between studies of high quality (score of 7 to 10) and low or moderate (score from 0 to 6). This cut-off point was chosen arbitrarily, since there are no stipulated classifications for this tool²⁰. In order to measure the agreement between the reviewers in assigning the scores for each study, an Intraclass Correlation Coefficient was performed via SPSS v. 20.0 software, which was classified as: poor (ICC <0.4), satisfactory (0.4 ≤ ICC <0.75) and excellent (ICC ≥ 0.75)²⁵.

Statistical Analysis

The data was initially analyzed by means of descriptive statistics, separated into subgroups according to the vertebral region associated with the changes and the instrument of analysis. A meta-analysis was performed using the Comprehensive Meta-Analysis, version 2.2.04 (Biostat, Inc. ©, Englewood, New Jersey) and MedCalc²⁶, version 11.0 (MedCalc Software, Mariakerke, Belgium) software by means of inferential statistics with the Higgins Inconsistency Test (I²)²¹ to verify homogeneity of the subjects. The heterogeneity was considered low if I²<50% and moderate / high if I²≥50%. In cases of I²≥50%, we opted for the adoption of the random models effect. The sensitivity analysis encompassed the exclusion of studies based on a sample calculation performed

for each subgroup, using the recommendations of Santos, Abbud and Abreu²⁷, that is, for each subgroup, sample calculations were performed from three different values of prevalence - low, intermediate and high (Table 2). Thus, the studies included in the systematic review that had not recruited the minimum sample to meet the proposed objective were excluded from the meta-analysis. However, even with the sensitivity analysis based on the inclusion of studies with a minimum sample size corresponding to the one obtained by the sample calculation, it was not possible to reduce or minimize the heterogeneity present in the studies. Therefore, the presentations of the meta-analysis in subgroups according to the vertebral region associated with the changes were maintained in general, in order to highlight the methodological differences between the studies. (Supplementary material)

Strength of evidence

In order to summarize the general quality of the evidence, the GRADE (Grading of Recommendations Assessment, Development, and Evaluation) system was used²⁸. GRADE classifies the quality of evidence and the strength of recommendation provided by systematic reviews, scientific advice, and clinical guidelines. It is a way of representing confidence in the information provided, by classifying the level of evidence and by expressing the emphasis so that a particular conduct is adopted or rejected in the case of reviews of clinical trials²⁹. We analyzed only observational studies, and only the classification of their level of evidence was used. The quality of evidence evaluation was based on the following criteria: design of studies included in the systematic review; methodological limitations of included studies;

Table 2: Studies used for sample calculation of each subgroup of analysis (cervical, thoracic, lumbar and scoliosis).

Analysis Subgroup	Article	Sample Size (n)	Prevalence (%)	Minimum sample size obtained by calculation (n)
Cervical	Santos et al. (2009)	247	11.7	162
	Lemos, Santos and Gaya (2012)	467	36.4	354
	Detsch and Candotti (2001)	154	66.2	345
Thoracic	Santos et al. (2009)	247	9.7	126
	Bastião et al. (2014)	420	40.5	369
	Noll et al. (2012)	65	66.1	345
Lumbar	Santos et al. (2009)	247	26.3	296
	Detsch and Candotti (2001)	154	31.2	329
	Pinho and Duarte (1995)	229	50.2	384
Scoliosis	Souza et al. (2013)	418	4.3	59
	Santo, Guimarães and Galera (2011)	210	18.1	227
	Sedrez et al. (2015)	59	49.1	384

Inconsistency (homogeneity of studies); whether the studies present direct evidence; accuracy of the results presented in included studies; and if the systematic review has a publication bias, not including all the published studies about the research problem. Using these criteria, the level of evidence was classified, among the four levels presented by the GRADE system: high quality, moderate quality, low quality and very low quality. In high quality evidences, it is very unlikely that additional research change the prevalence estimate presented by the systematic review; When the study presents very low quality evidence, its prevalence estimate is very uncertain²⁹, thus new studies are necessary.

RESULTS

The initial search identified 1193 studies, of which 221 were duplicates, thus 972 remained. However, 950 were excluded based on the title and abstract, so that only 22 remained for detailed analysis, six of which were excluded after screening for the eligibility criteria. Manual searches were carried out in the references of the 16 studies, and a further 12 studies were included. Thus, 29 studies were reviewed. Figure 1 shows the flowchart of the study selection, Table 3

summarizes the characteristics of the studies and Table 4 explains the methodological quality of the studies.

In order to summarize the evidence from the 27 studies, with regard to postural deviations in the spine, we can list, in ascending order of involvements: lumbar hyperlordosis (found in 14 studies, prevalence between 19% and 78.1%); Thoracic hyperkyphosis (found in 13 studies, prevalence between 9.7% and 49%) and scoliosis (in four studies, prevalence between 5.2% and 28%). In addition, the contrasting sample sizes between the studies, from 47 to 1340 students, stand out. The number of studies carried out in the different geographic regions of Brazil, conducted mainly in the South (n = 14), followed by the Southeast (n = 7), and in a smaller number in the Northeast (n = 4), also stand out, which is possibly related to regional economic constraints and resources.

Regarding the strength of evidence of this systematic review, based on the main criteria established by GRADE²⁹, 14 studies are in the high category of methodological quality, presenting an excellent agreement of punctuation between the reviewers (ICC = 0.833] 0.675; 0.918 [p <0.001), which implies a low risk of bias. However, with regard to inconsistency, high values were found, which reiterate the heterogeneity of the studies, making the quantitative informa-

Figure 1: Flowchart of the search and selection of studies according to PRISMA.




Table 3: Characteristics of the studies (in alphabetical order)

Article	Location	Participants	Methods	Positive events (n)
Baroni et al. (2015)36	Santa Cruz, RN	n = 212 (from 7 to 17 years) 58.5% female; 41.5% male	Visual inspection	Possible scoliosis = 123
Bastião et al. (2014)37	São Paulo State	n = 420 (from 1st to 8th grade elementary school) 60% female; 40% male	Photogrammetry	Cervical hyperlordosis = 87, Thoracic hyperkyphosis = 170, Lumbar hyperlordosis = 183
Bertolini, Gomes (1997)38	Maringá, PR	n = 200 (from 11 to 14 years) 59.5% female; 40.5% male	Visual inspection	Cervical hyperlordosis = 36, Thoracic hyperkyphosis = 38, Possible scoliosis = 29, Lumbar hyperlordosis = 53
Brianezi, Cajazeiro, Maifri-no (2011)39	Hortolândia, SP	n = 201 (from 7 to 10 years) 52.2% female; 47.8% male	Visual inspection	Lumbar hyperlordosis = 81, Thoracic hyperkyphosis = 88 Possible scoliosis = 102, Lumbar hyperlordosis = 99
Bueno, Rech (2013)40	Caxias do Sul, RS	n = 864 (from 8 to 15 years) 49% female; 51% male	Visual inspection	Thoracic rectification= 143, Thoracic hyperkyphosis=30 Possible scoliosis = 287, Lumbar hyperlordosis= 241 Lumbar rectification = 31, Curve inversion = 2
Contri, Petrucci, Perea (2009)41	Porto Ferreira, SP	n = 465 (from 7 to 12 years) 56% female; 44% male	Evaluation Form/ Visual	Thoracic hyperkyphosis = 117, Possible scoliosis = 64 Lumbar hyperlordosis = 144
Correa, Pereira, Silva (2005)42	Barra Mansa, RJ	72 (from 8 to 15 years)	Visual Inspection	Thoracic hyperkyphosis = 20, Possible scoliosis = 38 Lumbar hyperlordosis = 34
Detsch, Candotti (2001)43	Novo Hamburgo, RS	n = 154 (from 6 to 17 years) 100% female	Visual Inspection	Cervical anteriorization = 102, Thoracic hyperkyphosis = 16, Lumbar hyperlordosis = 48
Detsch et al. (2007)44	São Leopoldo, RS	n = 495 (from 14 to 18 years) 100% female	Visual Inspection	Possible scoliosis = 327
Dönert, Tomasi (2008)45	Pelotas, RS	n = 314 (from 9 to 16 years) 45.5% female; 54.5% male	Visual Inspection	Scoliosis = 28
Fornazari, Pereira (2008)46	Ribeirão Preto, SP	n = 497 47% female; 53% male	X-ray	Possible scoliosis = 108
Graup, Santos, Moro (2010)47	Florianópolis, SC	n = 288 (from 15 to 18 years) 46% female; 54% male	Visual Inspection	Lumbar hyperlordosis = 14, Lumbar rectification= 141
Lemos et al. (2005)48	General Câmara, RS	131 (from 10 to 13 years) 66.4% female; 36.6% male	Photogrammetry	Thoracic hyperkyphosis = 26, Lumbar hyperlordosis = 85 Lumbar rectification = 3
Lemos, Santos, Gaya (2012)49	Porto Alegre, RS	467 (from 10 to 16 years) 44.3% female; 55.7% male	Visual Inspection	Cervical rectification = 56, Cervical hyperlordosis = 114 Thoracic hyperkyphosis = 179
Martelli, Traebert (2006)50	Tangará, SC	344 (from 10 to 16 years) 52.9% female; 47.1% male	Visual Inspection	Lumbar hyperlordosis = 365, Thoracic hyperkyphosis = 38 Possible scoliosis = 30, Lumbar hyperlordosis = 70
Nery et al. (2010)51	Carlos Barbosa, RS	1340 (from 5th to 8th grade elementary school) 49% female; 51% male	Visual Inspection	Possible scoliosis = 19
Noll et al. (2012)52	Teutônia, RS	65 (from 11 to 16 years) 43% female; 57% male	Visual Inspection	Cervical anteriorization = 24 Cervical retroversion = 18 Thoracic hyperkyphosis = 26 Thoracic rectification= 17 Possible scoliosis = 41 Lumbar hyperlordosis = 30 Lumbar rectification = 2

Penha et al. (2005)53	São Paulo, SP	132 (from 7 to 10 years) 100% female	Photogrammetry	Possible scoliosis = 41, Lumbar hyperlordosis = 30 Lumbar rectification = 2, Thoracic hyperkyphosis = 45
Pereira et al. (2005)54	Jequié, BA	143 (from 10 to 15 years) 72.1% female; 27.9% male	Visual Inspection	Possible scoliosis = 60
Pinho, Duarte (1995)55	Florianópolis, SC	229 (from 7 to 10 years) 48.5% female; 51.5% male	Visual Inspection	Thoracic hyperkyphosis = 59, Possible scoliosis = 71 Lumbar hyperlordosis = 115
Rego, Scartoni (2008)56	Teresina, PI	47 34% female; 66% male	Visual Inspection	Possible scoliosis = 24
Rocha, Tamatsu, Vilela (2012)57	Quixadá, CE	228 (12 years) 64.5% female; 35.5% male	Visual Inspection	Possible scoliosis = 110
Rodrigues et al. (1985)58	Rio Grande, RS	135 (from 6 to 14 years) 54.9% female; 45.1% male	X-ray	Scoliosis = 7
Santo, Guimarães, Galera (2011)59	Cuiabá, MT	210 (3rd and 4th grades) 53.8% female; 46.2% male	X-ray	Scoliosis = 38, Cervical protusion = 29, Thoracic hyperkyphosis = 24, Possible scoliosis = 39, Lumbar hyperlordosis = 65
Santos et al. (2009)60	Jaguariúna, SP	247 (from 6 to 13 years) 47% female; 53% male	X-ray	Cervical protusion = 29, Thoracic hyperkyphosis = 24 Possible scoliosis = 39, Lumbar hyperlordosis = 65
Sedrez et al. (2015)61	Porto Alegre, RS	59 (from 7 to 18 years) 55.9% female; 44.1% male	X-ray	Thoracic hyperkyphosis = 30 Scoliosis = 29, Lumbar hyperlordosis = 19
Souza et al. (2013)62	Goiânia, GO	418 (from 10 to 14 years)	X-ray	Scoliosis = 18

Table 4: Results of methodological quality assessment and studies bias risk

Total Studies	Methodological quality and risk of bias										n of ✓
	1 st author (year) (Prevalence Critical Appraisal Instrument)										
1 st author (year)	1	2	3	4	5	6	7	8	9	10	
Baroniet al. (2015)36	✓	✓	✓	✓	✓	✓	?	✓	✓	✓	9
Bastião et al.(2014)37	?	✓	?	X	?	✓	✓	✓	✓	✓	6
Bertolini, Gomes (1997)38	?	?	?	X	?	✓	?	?	X	X	1
Brianezi, Cajazeiro and Maifrino (2011)39	?	?	?	X	?	✓	?	?	X	X	1
Bueno, Rech (2013)40	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	10
Contri, Petrucelli, Perea (2009)41	?	✓	?	X	✓	✓	?	✓	X	X	4
Correa et al. (2005)42	?	?	?	X	✓	✓	?	?	X	X	2
DetschandCandotti (2001)43	?	X	?	✓	?	✓	✓	✓	✓	✓	6
Detschet al. (2007)44	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	10
Döhnert, Tomasi (2008)45	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	10
Fornazari, Pereira (2008)46	?	✓	?	✓	X	✓	✓	✓	X	X	5
Graup, Santos, Moro (2010)47	✓	✓	✓	✓	?	✓	✓	✓	✓	✓	8
Lemos et al. (2005)48	?	?	?	✓	?	✓	✓	✓	X	X	4
Lemos, Santos e Gaya (2012)49	?	X	?	✓	?	✓	✓	✓	✓	✓	6
Martelli, Traebert (2006)50	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	10
Nery et al. (2010)51	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	10
Noll et al. (2012)52	✓	?	✓	✓	✓	✓	✓	✓	X	✓	8
Penha et al. (2005)53	?	✓	?	✓	?	✓	✓	✓	✓	✓	7
Pereira et al. (2005)54	✓	✓	?	✓	✓	✓	✓	✓	✓	✓	9
Pinho, Duarte (1995)55	?	?	?	✓	?	✓	?	✓	✓	✓	5
Rego, Scartoni (2008)56	?	?	?	✓	?	✓	?	?	X	X	3
Rocha, Tamatsu, Vilela (2012)57	?	✓	?	✓	?	✓	?	?	✓	✓	7
Rodrigues et al. (1985)58	?	?	?	✓	✓	✓	✓	✓	✓	✓	5
Santo, Guimarães and Galera (2011)59	✓	✓	?	?	✓	✓	✓	✓	✓	✓	9
Santos et al. (2009)63	✓	✓	✓	✓	?	✓	✓	✓	X	✓	8
Sedrez et al. (2014)64	✓	✓	✓	X	✓	✓	✓	✓	✓	✓	9
Souza et al. (2013)62	X	✓	X	✓	✓	✓	✓	✓	X	X	6

Responses to criteria: ✓ = Yes; X = No, ? = Unclear

1. Was the sample representative of the target population?
2. Were the participants adequately recruited?
3. Was the sample size adequate?
4. Have the individuals and the recruitment environment been described in detail?
5. Was the data analysis done with sufficient coverage of the identified sample?
6. Were standard objective criteria used to measure the condition?
7. Was the condition measured reliably?
8. Was the statistical analysis appropriate?
9. Have all major confounding factors/subgroups/differences been identified and considered?
10. Were subpopulations identified with objective criteria?

tion from analysis to meta-analysis inaccurate, a fact that is added to the high confidence intervals obtained in the calculations of prevalence by subgroup. Given the above, it is possible that future studies are likely to have a significant impact on the confidence in estimating the prevalence of postural deviations in the spine of schoolchildren, which makes this review with low strength of evidence.

■ DISCUSSION

The meta-analysis aimed to identify the prevalence of postural deviations in the spine of schoolchildren and adolescents in Brazil. Although the meta-analysis has a sample of individuals of over a thousand students in the analysis of each type of postural change, it was not possible to establish a consensus about the prevalence of anteroposterior and latero-lateral postural deviations in the spine of Brazilian schoolchildren. This result can be due the great heterogeneity of the studies, since they differ significantly regarding the methods used for evaluation of the posture and the sample space. Therefore, the prevalence of postural deviations were discrepant, reaching, thus, heterogeneity indexes above 90%.

Among the ways of assessing spinal posture, visual inspection depends exclusively on the experience of the professional performing the assessment, besides not allowing to quantify objectively the changes, being an evaluation associated with major errors and disagreements^{30,31}. Likewise, the postural evaluation by photogrammetry, although providing quantitative evidence on the curvatures of the spine³², is also subject to recurrence of errors, either due to the inherent difficulty of palpation or by the different mathematical procedures that the software have³³. In fact, the results of the studies that used photogrammetry were also discrepant with each other. Nevertheless, both methods, visual inspection and photogrammetry, were the most used by the studies included in this systematic review, factors that may contribute to the impossibility of reaching consensus on the prevalence of postural deviations in the Brazilian schoolchildren population.

Another factor that may have contributed to the variability of the results between the studies is a wide range of reference values for the classification of spinal curvatures. For example, for Bernhardt & Bridwell³⁴, Cobb angle values for a normal lumbar spine curvature ranged from 14° to 69°, while for Propst-Proctor and Bleck³⁵ they ranged from 22° to 54°. In this sense, depending on the reference values used in the studies, these different spectra may have generated large differences in the posture classifications of

the spine.

Despite the biases found in the methodologies of the studies included in this review, when the studies were submitted to the methodological quality and bias risk assessment through the Prevalence Critical Appraisal Instrument²⁴, 14 studies were classified with high quality. According to GRADE²⁹, this result indicates a low risk of bias in the systematic review. Nevertheless, the meta-analysis indicates an inconsistency, due to the high heterogeneity among the studies. In this perspective, it is emphasized that there are studies of high methodological quality presenting the panorama of the problem of postural deviations in Brazilian schoolchildren, however such panoramas cannot be extrapolated to the context of the country, as they show only the reality of the region in which each study was developed. An example of this question is the fact that there are students whose habit of sleeping in hammocks is so recurrent that this factor was evaluated in the study by Baroni *et al.*³⁶, being considered a protective factor for the development of scoliosis. This life habit was evaluated only in the study developed in north of Brazil, and was not found in any other Brazilian study. Therefore, since different regions of Brazil present different behavioral and life habits due to the different regional cultural influences, we can expect different types of postural deviations and different prevalence among their populations.

Thus, it was not possible to reach a consensus regarding the prevalence of postural deviations in the spine of Brazilian schoolchildren. From this perspective, we see the need for future studies designed to eliminate the biases pointed out by this systematic review, so that the Brazilian reality regarding the posture of the spine of schoolchildren can be documented. We understand that from a global knowledge, health promotion and education actions can be developed and put into practice at federal level, a fact that is still not possible today and therefore requires strategies at regional levels.

■ CONCLUSION

Based on the results obtained from the systematic review of articles developed in Brazil, we conclude that there is a low strength of evidence to establish a consensus about the values of prevalence of postural deviations in the spine of schoolchildren. We suggest that future studies be more rigorous in the screening and establishment of methodologies, as well as using validated instruments for evaluation, and investigating the macro and micro-regional differences of the states and the country.

■ REFERENCES

- Galvão CM, Sawada NO, Trevizan MA. Revisão sistemática: recurso que proporciona a incorporação das evidências na prática da enfermeira na enfermagem. *Rev Latino-Am Enferm*. 2004;12(3):549-56. DOI: <http://dx.doi.org/10.1590/S0104-11692004000300014>
- Skoffer B, Foldspang A. Physical activity and low-back pain in schoolchildren. *Eur Spine J*. 2008;17(3):373-9. DOI: <http://dx.doi.org/10.1007/s00586-007-0583-8>

3. Cakmak A, Yücel B, Özyalçın SN, Bayraktar B, Ural HI, Duruöz MT, *et al.* The frequency and associated factors of low back pain among a younger population in Turkey. *Spine (Phila Pa 1976)*. 2004;29(14):1567-72. DOI: <http://dx.doi.org/10.1097/01.BRS.0000131432.72531.96>
4. Paananen MV, Taimela SP, Auvinen JP, Tammelin TH, Kantomaa MT, Ebeling HE, *et al.* Risk factors for persistence of multiple musculoskeletal pains in adolescence : A 2-year follow-up study. *Eur J Pain*. 2010;14(10):1026-32. <http://dx.doi.org/10.1016/j.ejpain.2010.03.011>
5. Lis AM, Black KM, Korn H, Nordin M. Association between sitting and occupational LBP. *Eur Spine J*. 2007;16(2):283-98. DOI: <http://dx.doi.org/10.1007/s00586-006-0143-7>
6. Oshiro VA, Ferreira PG, Costa RF. Alterações posturais em escolares: uma revisão da literatura. *Rev Bras Ciênc Saúde*. 2007;5(13):15-22. DOI: <http://dx.doi.org/10.13037/rbcs.vol5n13.396>
7. Micheli L, Fehlandt AJ. Overuse injuries to tendons and apophyses in children and adolescents. *Clin Sports Med*. 1992;11(4):713-26.
8. Braccialli LMP, Vilarta R. Aspectos a serem considerados na elaboração de programas de prevenção e orientação de problemas posturais. *Rev Paul Educ Fís*. 2000;14(1):16-28.
9. Ainhagne M, Santhiago V. Cadeira e mochila escolares no processo de desenvolvimento da má postura e possíveis deformidades em crianças de 8-11 anos. *Colloq Vitae*. 2009;1(1):01-7. DOI: <http://dx.doi.org/10.5747/cv.2009.v01.n1.v001>
10. Grimmer K, Dansie B, Milanese S, Pirunsan U, Trott P. Adolescent standing postural response to backpack loads: a randomised controlled experimental study. *BMC Musculoskelet Disord*. 2002;3:10. DOI: <http://dx.doi.org/10.1186/1471-2474-3-10>
11. Politano RC. Levantamento dos desvios posturais em adolescentes de 11 a 15 anos em escola estadual do município de Cacoal - RO. Dissertação (Mestrado) - Universidade de Brasília. Brasília: 2006.
12. Aguiar NH, Bertolini SMMG. Estudo da incidência de cifose postural em crianças na faixa etária de 7 a 9 anos da rede escolar. *Arq Ciênc Saúde Unipar*. 1997;1(1):71-4.
13. Balagué F, Troussier B, Salminen JJ. Non-specific low back pain in children and adolescents: risk factors. *Eur Spine J*. 1999;8(6):429-38. DOI: <http://dx.doi.org/10.1007/s005860050201>
14. Silva MC, Fassa AG, Valle NCJ. Dor lombar crônica em uma população adulta do Sul do Brasil: prevalência e fatores associados. *Cad Saude Publica*. 2004;20(2):377-85. DOI: <http://dx.doi.org/10.1590/S0102-311X2004000200005>
15. Fathi M, Rezaei R. Comparison of height anomalies in middle school and high school students. *Teach Phy Edu*. 2011;11(1):46-53.
16. Zakeri Y, Baraz S, Gheibizadeh M, Nejad DB, Latifi SM. Prevalence of musculoskeletal disorders in primary school students in Abadan-Iran in 2014. *Int J Pediatr*. 2016;4(1):1215-23. DOI: <http://dx.doi.org/10.22038/ijp.2016.6265>
17. Murata Y, Utsumi T, Hanaoka E, Takahashi K, Yamagata M, Moriya H. Changes in lumbar lordosis in young patients with low back pain during a 10-year period. *J Orthop Sci*. 2002;7(6):618-22. DOI: <http://dx.doi.org/10.1007/s007760200111>
18. Sano A, Hirano T, Watanabe K, Endo N, Ito T, Tanabe N. Body mass index is associated with low back pain in childhood and adolescence: a birth cohort study with a 6-year follow-up in Niigata City , Japan. *Eur Spine J*. 2015; 24(3):474-81. DOI: <http://dx.doi.org/10.1007/s00586-014-3685-0>
19. Galvão TF, Pereira MG. Revisões sistemáticas da literatura: passos para sua elaboração. *Epidemiol Serv Saúde*. 2014;23(1):183-4. DOI: <http://dx.doi.org/10.5123/S1679-49742014000100018>
20. The Joanna Briggs Institute (JBI). Joanna Briggs Institute reviewers' manual 2014: the systematic review of prevalence and incidence data. Austrália: 2014; p.37.
21. Higgins J, Green S. *Cochrane handbook for systematic reviews of interventions*. [Cited 2016 Dec 15] Available from: <http://www.mri.gov.lk/assets/Uploads/Research/Cochrane-Hand-booktext.pdf>
22. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, *et al.* Meta-analysis of observational studies in epidemiology: a proposal for reporting. *JAMA*. 2000;283(15):2008-12. DOI: <http://dx.doi.org/10.1001/jama.283.15.2008>
23. Pereira MG, Galvão TF. Etapas de busca e seleção de artigos em revisões sistemáticas da literatura. *Epidemiol Serv Saúde*. 2014;23(2):369-71. DOI: <http://dx.doi.org/10.5123/S1679-49742014000200019>
24. Munn Z, Moola S, Riitano D, Lisy K. The development of a critical appraisal tool for use in systematic reviews addressing questions of prevalence. *Int J Heal Policy Manag*. 2014;3(3):123-8. DOI: <http://dx.doi.org/10.15171/ijhpm.2014.71>
25. Fleiss JN, Levin B, Paik MC. *Statistical methods for rates and proportions*. 3rd Ed. New York: 2004.
26. Schoonjans F. *Medcalc statistics for biomedical research: software manual*. Mariakerke: Medcalc Statistical Software; 2016; p. 295.
27. Santos GR, Abbud EL, Abreu AJ. Determination of the size of samples : an introduction for new researches. *Rev Científica Symposiun*. 2007;5(1):59-65.
28. Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, *et al.* GRADE guidelines : 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol*. 2011;64(4):383-94. DOI: <http://dx.doi.org/10.1016/j.jclinepi.2010.04.026>


29. Ministério da Saúde. Secretaria de Ciência, Tecnologia e Insumos Estratégicos Departamento de Ciência e Tecnologia. Diretrizes metodológicas: Sistema GRADE - Manual de graduação da qualidade de evidência e força de recomendação para tomada de decisão em saúde. Brasília: MS; 2014; p. 72.
30. Furlanetto TS, Chaise FDO, Candotti CT, Loss JF. Fidedignidade de um protocolo de avaliação postural. Rev Educ Física. 2011;22(3):411-9. DOI: <http://dx.doi.org/10.4025/reveducfis.v22i3.10124>
31. Chaise FO, Candotti CT, Torre ML, Furlanetto TS, Pelinson PPT, Loss JF. Validation, repeatability and reproducibility of a noninvasive instrument for measuring thoracic and lumbar curvature of the spine in the sagittal plane. Rev Bras Fisioter. 2011;15(6):511-7. DOI: <http://dx.doi.org/10.1590/S1413-35552011005000031>
32. D'Osualdo F, Schierano S, Cisotti C. The evaluation of the spine through the surface. The role of surface measurements in the evaluation and treatment of spine diseases in young patients. Eur Medicophys. 2002;38(3):147-52.
33. Iunes D, Bevilaqua-Grossi D, Oliveira A, Castro F, Salgado H. Análise comparativa entre avaliação postural visual e por fotogrametria computadorizada. Rev Bras Fisioter. 2009;13(4):308-15. DOI: <http://dx.doi.org/10.1590/S1413-35552009005000039>
34. Bernhardt M, Bridwell KH. Segmental analysis of the sagittal plane alignment of the normal thoracic and lumbar spines and thoracolumbar junction. Spine (Phila Pa 1976). 1989;14(7):717-21. DOI: <http://dx.doi.org/10.1097/00007632-198907000-00012>
35. Propst-Proctor SL, Bleck EE. Radiographic determination of lordosis and kyphosis in normal and scoliotic children. J Pediatr Orthop. 1983;3(3):344-6. DOI: <http://dx.doi.org/10.1097/01241398-198307000-00013>
36. Baroni MP, Sanchis GJ, de Assis SJ, dos Santos RG, Pereira SA, Sousa KG, *et al.* Factors associated with scoliosis in schoolchildren: a cross-sectional population-based study. J Epidemiol. 2015;25(3):212-20. DOI: <http://dx.doi.org/10.2188/jea.JE20140061>
37. Bastião MV, Carnaz L, Barbosa LF, Motta GC, Sato T de O. Posture and musculoskeletal pain in eutrophic, overweighted, and obese students. A cross-sectional study. Motriz: Rev Educ Fis. 2014;20(2):192-9. DOI: <http://dx.doi.org/10.1590/S1980-65742014000200009>
38. Bertolini SMMG, Gomes A. Estudo da incidência de cifose postural em adolescentes na faixa etária de 11 a 14 anos da rede escolar de Maringá. Rev Educ Física. 1997;8(1):105-10.
39. Brianezi L, Cajazeiro DC, Maifrino LBM. Prevalence of postural deviations in school of education and professional practice of physical education. J Morphol Sci. 2011;28(1):35-6.
40. Bueno RCS, Rech RR. Postural deviations of students in Southern Brazil. Rev Paul Pediatr. 2013;31(2):237-42. DOI: <http://dx.doi.org/10.1590/S0103-05822013000200016>
41. Contri DE, Petrucelli A, Perea DCBNM. Incidência de desvios posturais em escolares do 2o ao 5o ano do Ensino Fundamental. ConScientiae Saúde. 2009;8(2):219-24. DOI: <http://dx.doi.org/10.5585/conssaude.v8i2.1637>
42. Correa AL, Pereira JS, Silva MAG. Avaliação dos desvios posturais em escolares: estudo preliminar. Fisioter Bras. 2005;6(3):175-8.
43. Detsch C, Candotti CT. A incidência de desvios posturais em meninas de 6 a 17 anos da cidade de Novo Hamburgo. Movimento. 2001;7(15):43-56.
44. Detsch C, Luz AMH, Candotti CT, Oliveira DS, Lazzaron F, Guimarães LK, *et al.* Prevalência de alterações posturais em escolares do ensino médio em uma cidade no Sul do Brasil. Rev Panam Salud Pública. 2007;21(4):231-8. DOI: <http://dx.doi.org/10.1590/S1020-49892007000300006>
45. Dohnert MB, Tomasi E. Validade da fotogrametria computadorizada na detecção de escoliose idiopática adolescente. Rev Bras Fisioter. 2008;12(4):290-7. DOI: <http://dx.doi.org/10.1590/S1413-35552008000400007>
46. Fornazari LP, Pereira VCG. Prevalência de postura escoliótica em escolares do ensino fundamental. Cad Esc Saúde Fisioter. 2008;1(1):1-13.
47. Graup S, Santos SG, Moro ARP. Estudo descritivo de alterações posturais sagitais da coluna lombar em escolares da rede federal de ensino de Florianópolis. Rev Bras Ortop. 2010;45(5):453-9. DOI: <http://dx.doi.org/10.1590/S0102-36162010000500013>
48. Lemos A, Machado D, Moreira R, Torres L, Garlipp D, Lorenzi T, *et al.* Atitude postural de escolares de 10 a 13 anos de idade. Rev Perfil. 2005;7:53-9.
49. Lemos AT, Santos FR, Gaya ACA. Hiperlordose lombar em crianças e adolescentes de uma escola privada no Sul do Brasil : ocorrência e fatores associados. Cad Saude Publica. 2012;28(4):781-8. DOI: <http://dx.doi.org/10.1590/S0102-311X2012000400017>
50. Martelli RC, Traebert J. Estudo descritivo das alterações posturais de coluna vertebral em escolares de 10 a 16 anos de idade: Tangará-SC, 2004. Rev Bras Epidemiol. 2006;9(1):87-93. DOI: <http://dx.doi.org/10.1590/S1415-790X2006000100011>
51. Nery LS, Halpern R, Nery PC, Nehme KP, Stein AT. Prevalence of scoliosis among school students in a town in southern Brazil. Sao Paulo Med J. 2010;128(2):69-73. DOI: <http://dx.doi.org/10.1590/S1516-31802010000200005>
52. Noll M, Rosa BN, Candotti CT, Furlanetto TS, Gontijo KNS, Sedrez JA. Alterações posturais em escolares do en-

- sino fundamental de uma escola de Teutônia/RS. Rev Bras Ciênc Mov. 2012;20(2):32-42.
53. Penha PJ, João SMA, Casarotto RA, Amino CJ, Penteadó DC. Postural assessment of girls between 7 and 10 years of age. Clinics. 2005;60(1):9-16. DOI: <http://dx.doi.org/10.1590/S1807-59322005000100004>
54. Pereira LM, Barros PCC, Oliveira MND, Barbosa AR. Escoliose: triagem em escolares de 10 a 15 anos. Rev Saúde Com. 2005;1(2):134-43.
55. Pinho RA, Duarte MFS. Análise postural em escolares de Florianópolis - SC. Rev Bras Ativ Física Saúde. 1995;1(2):49-58.
56. Rego ARON, Scartoni FR. Alterações posturais de alunos de 5ª e 6ª séries do ensino fundamental. Fit Perform J. 2008;7(1):10-15. DOI: <http://dx.doi.org/10.3900/fpj.7.1.10.p>
57. Rocha JCT, Tatmatsu DIB, Vilela DA. Associação entre uso de mochilas escolares e escoliose em adolescentes de escolas públicas e privadas. Motricidade. 2012;8(S2):803-9.
58. Rodrigues CAC, Teixeira MAM, Casartelli MRO. Escoliose: levantamento epidemiológico em alunos da escola estadual Lilia Neves. Vittalle. 1985;1:67-76.
59. Espírito Santo A, Guimarães LV, Galera MF. Prevalência de escoliose idiopática e variáveis associadas em escolares do ensino fundamental de escolas municipais de Cuiabá, MT, 2002. Rev Bras Epidemiol. 2011;14(2):347-56.
60. Santos CIS, Cunha ABN, Braga VP, Saad IAB, Ribeiro MAGO, Conti PBM, *et al.* Ocorrência de desvios posturais em escolares do ensino público fundamental de Jaguariúna, São Paulo. Rev Paul Pediatr. 2009;27(1):74-80. DOI: <http://dx.doi.org/10.1590/S0103-05822009000100012>
61. Sedrez JA, Rosa MIZ, Noll M, Medeiros FS, Candotti CT. Risk factors associated with structural postural changes in the spinal column of children and adolescents. Rev Paul Pediatr. 2015;33(1):72-81. DOI: <http://dx.doi.org/10.1016/j.rpped.2014.11.012>
62. Souza FI, Ferreira RB, Labres D, Elias R, Souza APM, Pereira RE. Epidemiologia da Escoliose Idiopática do Adolescente em alunos da rede pública de Goiânia-GO. Acta Ortop Bras. 2013;21(4):2012-4. DOI: <http://dx.doi.org/10.1590/S1413-78522013000400008>
63. Santos MM, Silva MPC, Sanada LS, Alves CRJ. Análise postural fotogramétrica de crianças saudáveis de 7 a 10 anos: confiabilidade interexaminadores. Rev Bras Fisioter. 2009;13(4):350-5. DOI: <http://dx.doi.org/10.1590/S1413-35552009005000047>
64. Sedrez JA, Furlanetto TS, Noll M, Gontijo KNS, Rosa BN, Candotti CT. Relação entre alterações posturais e fatores associados em escolares do ensino fundamental. Rev Baiana Saúde Pública. 2014;38(2):279-96. DOI: <http://dx.doi.org/10.5327/Z0100-0233-2014380200005>

Supplementary material

Table 1: Results of the meta-analysis for prevalence of alterations in the cervical spine


Article	Sample Size	Proportion (%)	95% CI	Weight (%)
Bastião et al. (2014)	420	20.7	16.9 – 24.9	23.9
Lemos, Santos and Gaya (2012)	467	36.4	32 – 40.9	26.6
Brianezi, Cajazeiro and Maifrino (2011)	201	40.3	33.4 – 47.4	11.5
Bertolini and Gomes (1997)	200	18	12.9 – 24	11.4
Detsch and Candotti (2001)	154	66.2	58.2 – 73.6	8.8
Noll et al. (2012)	65	64.6	51.8 – 76.1	3.7
Santos et al. (2009)	247	11.7	8 – 16.4	14.1
Total (fixed effects)	1754	30.2	28.1 – 32.5	100
Total (random effects)	1754	35.3	22.2 – 49.7	100



Heterogeneity test: $Q = 221.17$; $DF = 6$; $p < 0.001$; $I^2 = 97.3\%$; 95% CI for $I^2 = 95.9 - 98.2$

Table 2: Results of the meta-analysis for prevalence of changes in the thoracic spine


Article	Sample Size	Proportion (%)	95% CI	Weight (%)
Sedrez et al. (2015)	59	50.8	37.5 – 64.1	1.5
Bastião et al. (2014)	420	40.5	35.7 – 45.3	10.3
Brianezi, Cajazeiro and Maifrino (2011)	201	43.8	36.8 – 50.9	5
Santos et al. (2009)	247	9.7	6.3 – 14.1	6.1
Martelli and Traebert (2006)	344	11	7.9 – 14.8	8.5
Detsch and Candotti (2001)	154	10.4	6 – 16.3	3.8
Lemos et al. (2005)	131	19.8	13.4 – 27.7	3.2
Bueno and Rech (2013)	854	20	17.4 – 22.8	21
Pinho and Duarte (1995)	229	25.8	20.2 – 31.9	6.7
Bertolini and Gomes (1997)	200	19	13.8 – 25.1	4.5
Correa, Pereira and Silva (2005)	72	27.8	17.9 – 39.6	1.8
Lemos, Santos and Gaya (2012)	467	38.3	33.9 – 42.9	11.5
Noll et al. (2012)	65	66.1	53.3 – 77.4	1.6
Penha et al. (2005)	132	34	26.1 – 42.8	3
Contri, Petrucelli and Perea (2009)	465	25	21.3 – 29.4	11.5
Total (fixed effects)	4050	25.5	24.2 – 26.9	100
Total (random effects)	4050	28	21.5 – 35	100



Heterogeneity test: $Q = 316$; $DF = 14$; $p < 0.001$; $I^2 = 95.6\%$; 95% CI for $I^2 = 93.9 - 96.7$

Table 3: Results of the meta-analysis for prevalence of lumbar spine alterations


Article	Sample Size	Proportion (%)	95% CI	Weight (%)
Bastião et al. (2014)	420	43.5	38.8 – 48.5	15.4
Brianezi, Cajazeiro and Maifrino (2011)	201	49.2	42.1 – 56.4	7.4
Bueno and Rech (2013)	864	27.9	24.9 – 31	31.6
Santos et al. (2009)	247	26.3	20.9 – 32.3	9
Pinho and Duarte (1995)	229	50.2	43.3 – 56.9	8.4
Bertolini and Gomes (1997)	200	26.5	20.5 – 33.2	7.3
Correa, Pereira and Silva (2005)	72	47.2	35.3 – 59.3	2.7
Martelli and Traebert (2006)	344	20.3	16.2 – 25	12.6
Detsch and Candotti (2001)	154	31.2	23.9 – 39.1	5.6
Total (fixed effects)	2731	32.9	31.2 – 34.7	100
Total (random effects)	2731	35.7	28.1 – 42.8	100



Heterogeneity test: $Q = 124.9$; $DF = 8$; $p < 0.001$; $I^2 = 93.6\%$; 95% CI for $I^2 = 89.9 - 95.2$

Table 4: Results of the meta-analysis for prevalence of Scoliosis

Article	Sample Size	Proportion (%)	95% CI	Weight (%)
Sedrez et al. (2015)	59	49.1	35.9 – 62.5	3.6
Santo, Guimarães and Galera (2011)	210	18.1	13.1 – 23.9	12.9
Döhnert and Tomasi (2008)	314	8.9	6 – 12.6	19.3
Souza et al. (2013)	418	4.3	2.6 – 6.7	25.5
Rodrigues et al. (1985)	135	5.2	2.1 – 10.4	8.3
Fornazari and Pereira (2008)	497	8.8	6.5 – 11.7	30.4
Total (fixed effects)	1633	9.3	7.9 – 10.8	100
Total (random effects)	1633	12.9	6.7 – 20.8	100



Heterogeneity test: $Q = 86.12$; $DF = 5$; $p < 0.001$; $I^2 = 94.2\%$; 95% CI for $I^2 = 89.9 - 96.6$

This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Resumo

Introdução: A Era da informação e da tecnologia tem causado grande impacto na vida dos cidadãos. As invenções de máquinas, de automóveis, da televisão e do computador, induziram os indivíduos a adotarem a postura corporal “sentada” a fim de se adaptar as novas necessidades tecnológicas.

Objetivo: Estimar a prevalência de alterações posturais na coluna vertebral de escolares brasileiros.

Método: Foram realizadas buscas nas bases de dados EMBASE, LILACS, PubMed, SCOPUS, SciELO, Science Direct, e Web of Science, além de buscas manuais a fim de identificar estudos que avaliassem a prevalência de alterações posturais na coluna vertebral em escolares brasileiros. Dois revisores independentes realizaram a seleção dos estudos, avaliaram a qualidade metodológica e o risco de viés dos estudos selecionados e extraíram os dados. Foi realizada a análise de homogeneidade interestudos e a qualidade do nível de evidência foi avaliada utilizando o sistema GRADE.

Resultados: Foram incluídos 29 estudos, dos quais foram extraídas as frequências de eventos positivos para as alterações na coluna cervical, torácica, lombar, bem como a frequência de escoliose entre os escolares. Mesmo realizando a metanálise separada por subgrupos de acordo com a região vertebral avaliada, o nível de heterogeneidade ficou a cima dos 90%, não sendo possível estipular a prevalência de alterações posturais na coluna vertebral em escolares brasileiros a partir da metanálise.

Conclusão: Existe baixa força de evidência para se estabelecer um consenso acerca dos valores de prevalência de desvios posturais na coluna vertebral de escolares brasileiros.

Palavras-chave: postura, coluna, criança, adolescente.