

## HEART RATE VARIABILITY IN METABOLIC SYNDROME

Rodrigo Daminello Raimundo<sup>1</sup>, John J. Godleski<sup>1</sup>

DOI: <http://dx.doi.org/10.7322/jhgd.96757>

Metabolic syndrome (MetS) is a cluster of metabolic risk factors for cardiovascular disease and type 2 diabetes. It is also associated with insulin resistance (glucose intolerance) and obesity or overweight<sup>1,2</sup>. According to the International Diabetes Federation, a quarter of the world's adult population has MetS, and more cases are diagnosed in males over 40 years of age<sup>1</sup>.

MetS occurs when a person has three or more of the following measurements: (1) Abdominal obesity - waist circumference of 40 inches or more in men, and 35 inches or more in women; (2) Triglyceride level of 150 milligrams per deciliter of blood (mg/dL) or more; HDL cholesterol of less than 40 mg/dL in men or less than 50 mg/dL in women; (3) Systolic blood pressure of 130mmHg or greater, or diastolic blood pressure of 85mmHg or greater; and (4) Fasting glucose of 100 mg/dL or greater<sup>1,3</sup>.

There are increased risk for: Atherosclerosis, Peripheral vascular disease, Coronary heart disease, Heart attack, Stroke and Type 2 diabetes. *The National Cholesterol Education Program's Adult Treatment Panel III report* (ATP III) identified the proinflammatory state with elevations of C-reactive protein and a prothrombotic state with increased plasma plasminogen activator inhibitor and fibrinogen as associated with the metabolic syndrome<sup>4</sup>.

Many of the factors that contribute to metabolic syndrome can be addressed through lifestyle changes: balanced diet (maintaining a healthy weight, reduce fat intake, change the

consumption of saturated fats to unsaturated fats, increase the intake of fruits, vegetables, legumes and whole grains and reduced intake of sugar) and exercise (aerobic and muscle strengthening). The World Health Organization recommends 30 minutes of activity at least five times a week<sup>1,2,4,5</sup>.

Promising results of an important study published in the *BMC Public Health*<sup>6</sup> was *The Study Cardiovascular Risk in Adolescents* entitled "ERICA". This study aimed to estimate the prevalence of cardiovascular risk factors including metabolic syndrome in adolescents aged 12 to 17 years in Brazilian cities with over 100,000 inhabitants. We collected about 85,000 adolescents. The participants answered a questionnaire on demographic characteristics and lifestyle (physical activity, smoking, alcohol), mental disorders, reproductive and oral health, and blood samples for fasting glucose, total cholesterol, HDL-cholesterol, LDL - cholesterol, triglycerides, and glycated hemoglobin. It is expected that with the data from "ERICA" able to contribute to the knowledge about risk factors for atherosclerosis, obesity, diabetes mellitus among adolescents and serve to support the development of health policies

ATP III recommended first-line therapy the increased physical activity. The studies with exercise have increased in the scientific literature, between these studies autonomic modulation in patients with metabolic syndrome and exercise have risen too<sup>5,7</sup>. The heart autonomic modulation can be measured noninvasively, this process is known the analysis of heart rate variability (HRV)<sup>8</sup>. HRV can be used to

<sup>1</sup> Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, MA.  
**Corresponding author:** rodaminello@yahoo.com.br

**Suggested citation:** Raimundo RD, Godleski JJ. Heart rate variability in metabolic syndrome. *Journal of Human Growth and Development*. 2015; 25(1): 7-10.

Manuscript submitted February 16 2015, accepted for publication February 23 2015.

estimate autonomic nervous control of the cardiovascular system and reduced HRV significantly increases cardiovascular mortality<sup>9</sup>.

There is strong evidence that obesity leads to increased sympathetic modulation and a decreased vagal tone leading to a lack of autonomic control (characterized by reduction of heart rate variability) that can bring about changes in the renin-angiotensin-aldosterone system in alpha agonists and central alpha 2-1 and beta-adrenergic receptors. In addition, changes in plasma catecholamines and increased muscle sympathetic nerve activity are important characteristics in obese individuals<sup>8,9,10</sup>. All these factors contribute further to increased blood pressure<sup>11,12,13</sup>.

Souza et al<sup>14</sup> have reviewed studies of HRV in obese children, and find that obese children have lower activity of the parasympathetic nervous system. In addition,<sup>8-13</sup> there is evidence showing that interventions such as diet and physical activity can promote improvements in autonomic nervous system balance in this population.

Vanderlei et al<sup>15,16,17</sup> showed autonomic dysfunction characterized by reduced sympathetic modulation in obese children. The RRTri, TINN, SDNN, RMSSD, pNN50, SD1, SD2, LF and HF indices were lower among the obese children when compared to the eutrophic group.

Vanderlei, Vanderlei and Garner<sup>18</sup>, in an elegant study in this issue of JGDH publish important contribution in the area of HRV. This is the article "Heart rate dynamics by novel chaotic global to HRV in obese Youths" which is innovative evaluation of obese youth by chaotic behavior of HRV. Time heart rate series can be studied by means of statistical methods, this article studies the dynamic point of view, combining the nonlinear deterministic systems (chaotic behavior). Eighty-six young people were distributed in two groups (obese and control). The results illustrate that there is great variation in both the mean values and standard deviation for both young non-obese and obese, there is an increase in the chaotic response when going from non-obese to young obese, thus showing that the youth obesity increases the chaotic response.

Stuckey et al<sup>10</sup> systematically reviewed associations between heart rate variability and metabolic syndrome. Fourteen studies were included. Heart rate variability generally was reduced

in women with metabolic syndrome compared to those without, while results in men were inconsistent. Time and frequency domain heart rate variability parameters were associated with individual metabolic syndrome risk factors, though sex differences exist. The authors concluded HRV is altered differently in men and women with metabolic syndrome.

Stuckey et al<sup>19</sup> evaluated whether an 8-week exercise intervention would improve MetS risk factors and HRV in a population with MetS risk factors. The waist circumference and diastolic BP were reduced and maximal oxygen uptake (VO<sub>2</sub>max) increased. Low frequency (LF) was reduced and high frequency (HF) was increased. Thus, exercise intervention improved MetS risk factors and HRV parameters, but only changes in systolic BP were associated with improved autonomic function.

Koskinen et al<sup>20</sup> associated the presence of the MetS with lower HF and LF in men and women, and with higher LF/HF ratio in women. In young men, waist circumference was the strongest individual MetS component that associated with HRV. After adjustments for age and heart rate, MetS was associated with lower HF and higher LF/HF ratio in women.

Jarczok et al<sup>21</sup> recruited 2441 study participants and affirmed a negative correlation between HRV and glycemic status that appeared to be almost linear in a large cohort of healthy workers. Liao et al<sup>22</sup> examined the association between the level of cardiac autonomic activity and MetS disorders. HRV indices were significantly lower in individuals with MetS. An increase in fasting insulin was associated with 88% higher odds of having a lower HF. The pattern of associations was similar for LF and SDNN.

In recent years HRV has been an important instrument to evaluate autonomic modulation. The association of HRV indices with appropriate forms of exercise prescription can raise awareness for better treatment of patients with metabolic syndrome. MetS presented alterations in the autonomic nervous system, characterized by decreases in parasympathetic activity and global HRV, but the reduction in sympathetic modulation is not entirely clear. HRV analysis by chaotic behavior can bring an advance for evaluation and treatment of obese, diabetic, hypertensive and metabolic syndrome.

## REFERENCES:

1. Alberti KG, Zimmet P, Shaw J. Metabolic syndrome – a new world-wide definition. A Consensus Statement from the International Diabetes Federation. *Diabet Med.* 2006; 23(5): 469-80. 23(5): 469-80. Doi: 10.1111/j.1464-5491.2006.01858.x.
2. World Health Organization. Definition, diagnosis and classification of diabetes mellitus and its complications: report of a WHO consultation. Geneva, Switzerland: World Health Organization. Available from: [http://whqlibdoc.who.int/hq/1999/WHO\\_NCD\\_NCS\\_99.2.pdf](http://whqlibdoc.who.int/hq/1999/WHO_NCD_NCS_99.2.pdf).
3. Lopes HF, Egan BM. Autonomic Dysregulation and the Metabolic Syndrome: Pathologic partners in an emerging global pandemic. *Arq Bras Cardiol.* 2006; 87(4): 538-547.
4. Grundy SM, Brewer HB Jr, Cleeman JI, Smith SC Jr, Lenfant C; National Heart, Lung, and Blood Institute; American Heart Association. Definition of metabolic syndrome: report of the National Heart, Lung, and Blood Institute/American Heart Association conference on scientific related to definition. *Arterioscler Thromb Vasc Biol.* 2004 Feb; 24(2): e13-8. Doi: 10.1161/01.ATV.0000111245.75752.C6.
5. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA.* 2001; 285(19): 2486-97.
6. Bloch KV, Szklo M, Kuschner MCC, Abreu GA, Barufaldi LA, Klein CH. The study of cardiovascular risk in adolescents – ERICA: rationale, design and sample characteristics of a national survey examining cardiovascular risk factor profile in Brazilian adolescents. *BMC Public Health.* 2015; 15:94. Doi: 10.1186/s12889-015-1442-x. Page 8.
7. Lorenzo C, Serrano-Ríos M, Martínez- Lorenzo C, Serrano-Ríos M, Martínez-Larrad MT, González-Sánchez JL, Seclén S, Villena A, et al. Geographic variations of the International Diabetes Federation and the National Cholesterol Education Program Adult Treatment Panel III definitions of the metabolic syndrome in nondiabetic subjects. *Diabetes Care.* 2006 Mar; 29(3): 685-91. Doi: 10.2337/diacare.29.03.06.dc05-1796.
8. Grassi G, Seravalle G, Colombo M, Bolla G, Cattaneo BM, Cavagnini F, et al. Body weight reduction, sympathetic nerve activity, and arterial baroreflex in obese normotensive humans. *Circulation.* 1998 May; 97(20): 2037-42.
9. Abreu LC. Heart rate variability as a functional marker of development. *J Hum Growth Dev.* 2012; 22(3): 279-281.
10. Stuckey MI, Tulppo MP, Kiviniemi AM, Petrella RJ. Heart rate variability and the metabolic syndrome: a systematic review of the literature. *Diabetes Metab Res Rev.* 2014 Nov; 30(8):784-93. doi: 10.1002/dmrr.2555.
11. Carnethon MR, Prineas RJ, Tempresa M, Zhang ZM, Uwaifo G, Molitch ME. The association among autonomic nervous system function, incident diabetes, and intervention arm in the Diabetes Prevention Program. *Diabetes Care.* 2006 Apr; 29(4):914-9.
12. La Rovere MT, Bigger JT Jr, Marcus FI, Mortara A, Schwartz PJ. Baroreflex sensitivity and heart rate variability in prediction of total cardiac mortality after myocardial infarction. ATRAMI (autonomic tone and reflexes after myocardial infarction) investigators. *Lancet.* 1998; 351(9101): 478-484.
13. Kleiger RE, Miller JP, Bigger JT Jr, Moss AJ. Decreased heart rate variability and its association with increased mortality after acute myocardial infarction. *Am J Cardiol.* 1987; 59(4): 256-262.
14. Souza NM, Rossi RC, Vanderlei FM, Vitor ALRV, Bernardo AFB, Goncalves ACCR, et al. Heart rate variability in obese children. *J Hum Growth Dev.* 2012; 22(3):328-333.
15. Vanderlei LC, Pastre CM, Freitas Júnior IF, Godoy MF. Geometric indexes of heart rate variability in obese and eutrophic children. *Arq Bras Cardiol.* 2010; 95(1): 35-40. Doi: <http://dx.doi.org/10.1590/S0066-782X2010005000082>
16. Vanderlei LC, Pastre CM, Freitas Júnior IF, Godoy MF. Analysis of cardiac autonomic modulation in obese and eutrophic children. *Clinics.* 2010; 65(8): 789-92. Doi: <http://dx.doi.org/10.1590/S1807-59322010000800008>
17. Vanderlei LCM, Pastre CM, Júnior IFF, de Godoy MF. Fractal correlation of heart rate variability in obese children. *Auton Neurosci.* 2010; 155(12): 125-9. Doi: 10.1016/j.autneu.2010.02.002
18. Vanderlei FM, Vanderlei LCMV, Garner DM. Heart rate dynamics by novel chaotic globals to HRV in obese youths. *J Human Growth Dev.* 2015; 25(1): 11-17.
19. Stuckey MI, Kiviniemi AM, Petrella RJ. Diabetes and technology for increased activity study: the effects of exercise and technology on heart rate variability and metabolic syndrome risk factors.

- Front Endocrinol (Lausanne). 2013; 19(4): 121. Doi: 10.3389/fendo.2013.00121
20. Koskinen T, Kähönen M, Jula A, Mattsson N, Laitinen T, Keltikangas-Järvinen L, et al. Metabolic syndrome and short-term heart rate variability in young adults. The cardiovascular risk in young Finns study. *Diabet Med.* 2009; 26(4): 354-61. Doi: 10.1111/j.1464- 5491.2009.02686.x.
21. Jarczok MN, Li J, Mauss D, Fischer JE, Thayer JF. Heart rate variability is associated with glycemic status after controlling for components of the metabolic syndrome. *Int J Cardiol.* 2013; 167 (3): 855-61. Doi: 10.1016/j.ijcard.2012.02.002.
22. Liao D1, Sloan RP, Cascio WE, Folsom AR, Liese AD, Evans GW, et al. Multiple metabolic syndrome is associated with lower heart. The Atherosclerosis Risk in Communities Study. *Diabetes Care.* 1998; 21(12): 2116-22.