

## The moduli of Certain Curves of Genus Three in Characteristic Two

**Jaime E. A. Rodriguez**

Departamento de Matemática, FEIS, UNESP  
Av. Brasil, Centro 56, Cx P. 31, 15385-000, Ilha Solteira, SP, Brazil.  
*E-mail address:* `jaime@mat.feis.unesp.br`

**Oscar Paz la Torre**

Universidade Estadual do Norte Fluminense Darcy Ribeiro, LCMAT  
Av. Alberto Lamego 2000, CEP: 28013-600, Campos dos Goytacazes, Rio de Janeiro, Brazil.  
*E-mail address:* `oscar@uenf.br`

**Renato Vidal Martins**

Departamento de Matemática, ICEx, UFMG  
Av. Antônio Carlos 6627, 30123-970, Belo Horizonte MG, Brazil  
*E-mail address:* `renato@mat.ufmg.br`

**Abstract.** We study curves of genus 3 over algebraically closed fields of characteristic 2 with the canonical theta characteristic totally supported in one point. We compute the moduli dimension of such curves and focus on some of them which have two Weierstrass points with Weierstrass directions towards the support of the theta characteristic. We answer questions related to order sequence and Weierstrass weight of Weierstrass points.

*Key words:* curves of genus three, characteristic two, theta characteristic.

### 1. Introduction

In this article we study curves of genus 3 over fields of characteristic 2. In the case when the ground field is finite we have a recent wide classification (cf. [NR]). Here we assume the ground field is algebraically closed and we also assume the canonical theta characteristic is totally supported in one point.

So if  $C$  is such a curve then we can find a canonical divisor on  $C$  of the form  $4P_0$  where  $P_0$  is precisely the support of the theta characteristic. These curves must be non-hyperelliptic and hence canonically embedded in  $\mathbb{P}^2$  as a plane quartic described by an affine equation in a way that  $P_0$  is the origin. We have the following result.

**Theorem 1.** *The isomorphism classes of irreducible nonsingular projective curves of genus 3 over an algebraically closed field of characteristic 2, with the canonical theta characteristic totally supported at one point, form an algebraic variety of dimension 4.*

## 2. Preliminaries

Let  $C$  be an irreducible non-singular algebraic curve of genus 3 over a field of characteristic 2. And let us also assume that there exists a canonical theta characteristic  $\theta_0 = |\frac{1}{2}div(dx)|$  on  $C$  (with  $x$  a separating variable) which can be represented by a divisor of the form  $2P_0$  for a certain  $P_0 \in C$ . In this situation  $4P_0$  is a positive canonical divisor so that the point  $P_0$  has canonical order sequence 0, 1, 4. It follows that the curve is nonhyperelliptic and hence  $h^0(\mathcal{O}_C(2P_0)) = 1$  and  $2P_0$  is the only positive divisor in the class  $\theta_0$ .

Since  $C$  is nonhyperelliptic let us consider it canonically embedded in  $\mathbb{P}^2$  as a smooth plane quartic given by

$$f(x, y) = \sum_{i+j \leq 4} c_{ij} x^i y^j, \quad c_{ij} \in k. \quad (1)$$

For a general plane curve of genus  $g$  over a field of characteristic  $p$  we have that the *Cartier operator*

$$\mathcal{C} : \Omega_1(0) \longrightarrow \Omega_1(0)$$

which acts on the space of regular differentials of  $C$  can be expressed as

$$\mathcal{C}(hdx) = - \left( \frac{d^{p-1}h}{dx^{p-1}} \right)^{1/p} dx$$

(cf. [SV2]). The *Hasse-Witt invariant*  $\sigma$  is defined as the rank of the matrix

$$(h_{ij})(h_{ij}^p) \cdots (h_{ij}^{p^{g-1}})$$

for  $(h_{ij})$  the *Hasse-Witt matrix*, that is,  $(h_{ij}^{1/p})$  represents the Cartier operator.

In general a curve  $C$  of genus 3 admits a number of 7, 4, 2 or 1 bitangents,

depending on the values  $\sigma = 3, 2, 1$  or  $0$  of the Hasse-Witt invariant (cf. [SV2] pg. 60). In our case, as the canonical theta characteristic  $\theta_0$  does have a section, the Cartier operator necessarily has a non-trivial kernel, and so  $\sigma = 2, 1$  or  $0$ .

In the case of a non-singular plane curve (such as our canonical plane quartic) given by  $f(x, y) = 0$  as in (1), the differential  $\omega = \frac{dx}{f_y}$  is regular and

$$H^0(\mathcal{O}_C(\text{div}(\omega))) = \{ h \in k[x, y] \mid \deg(h) \leq \deg(f) - 3 \}.$$

We have that ([SV2] Theorem 1.1) in the case of characteristic 2 yields the formula for the Cartier operator

$$\mathcal{C}(h\omega) = \left( \frac{\partial^2}{\partial x \partial y} h f \right)^{1/2} \omega,$$

giving the Hasse-Witt matrix

$$H = \begin{pmatrix} c_{11} & c_{01} & c_{10} \\ c_{31} & c_{21} & c_{30} \\ c_{13} & c_{03} & c_{12} \end{pmatrix}.$$

We will use the theory of Weierstrass points, for which we refer to ([SV1] Section 1). In the case of a non-singular plane curve the results we need are collected below.

To compute the (canonical) Weierstrass points in curves of genus 3 over a field of characteristic 2 we use the classical *Wronskian* since from [K] there are no non-classical curves in this situation. Using the separating variable  $x$  we obtain

$$W_x^{0,1,2} = \det \begin{pmatrix} 1 & x & y \\ 0 & 1 & D_x^{(1)}(y) \\ 0 & 0 & D_x^{(2)}(y) \end{pmatrix} = D_x^{(2)}(y),$$

where  $D_x^{(i)}(y)$  stands for the  $i^{\text{th}}$  Hasse-Schmidt derivative. To compute the Wronskian we take *generic Taylor expansions*

$$\mathcal{T}(x) = x + t$$

$$\mathcal{T}(y) = y + D_x^{(1)}(y)t + D_x^{(2)}(y)t^2 + \dots$$

and use that

$$f(\mathcal{T}(x), \mathcal{T}(y)) = \sum_{i,j} c_{ij} \mathcal{T}(x)^i \mathcal{T}(y)^j = 0.$$

This yields

$$D_x^{(2)}(y) = \frac{w_K}{f_y^3},$$

where the numerator  $w_K$  is given by

$$w_K = f_x f_y \sum_{i,j \equiv 1 \pmod 2} c_{ij} x^{i-1} y^{j-1} + f_y^2 \sum_{i \equiv 2,3 \pmod 4} c_{ij} x^{i-2} y^j + f_x^2 \sum_{j \equiv 2,3 \pmod 4} c_{ij} x^i y^{j-2},$$

for

$$f_x = \sum_{i \equiv 1 \pmod 2} c_{ij} x^{i-1} y^j \quad \text{and} \quad f_y = \sum_{j \equiv 1 \pmod 2} c_{ij} x^i y^{j-1}.$$

The *ramification divisor* (cf. [SV1] pg. 3) is given by

$$\begin{aligned} \mathcal{R}_K &= \operatorname{div} (D_x^{(2)}(y)) + 3 \operatorname{div} (dx) + 3E \\ &= \operatorname{div} (w_K) + 3 \operatorname{div} \left( \frac{dx}{f_y} \right) + 3E \\ &= \operatorname{div} (w_K) + 6E \end{aligned}$$

where  $E = \operatorname{div} (\omega) = \operatorname{div} \left( \frac{dx}{f_y} \right)$  is the intersection divisor of the curve with the infinite line. The finite Weierstrass points are thus the zeros (counted with multiplicities) of the numerator  $w_K$  of  $D_x^{(2)}(y)$ . In our case of genus 3 and characteristic 2 there are altogether 24 Weierstrass points.

If  $P$  is a Weierstrass point then its order sequence may be only 0, 1, 3 or 0, 1, 4 and as a consequence of ([SV1] Theorem 1.5) its Weierstrass weight is 1 in the first case and greater than 2 in the second. The intersection divisor of the tangent of the curve at  $P$  is then  $3P + Q$  (with  $P = Q$  if the order sequence at  $P$  is 0, 1, 4). If  $Q$  itself is a Weierstrass point then we will say that  $P$  has a *Weierstrass direction towards  $Q$*  and

$$P \xrightarrow{w} Q$$

will denote this situation.

### 3. The moduli problem

We start this section with a result used to rule out some (in principle) possible degenerate situations. Recall that there are curves in prime characteristics having just one Weierstrass point.

**Proposition 3.1.** *The Weierstrass points of a smooth plane quartic in characteristic 2 with the canonical theta characteristic supported at one point are non collinear.*

*Proof.* If all Weierstrass points were collinear then we can suppose that the line containing them is the infinite line. In this case the numerator  $w_K$  of the Wronskian would be a non-zero constant, as there will be no finite Weierstrass points.

We can further assume that the canonical theta characteristic  $\theta_0$  contains the divisor  $2P_0$ , for  $P_0 = (1 : 0 : 0)$ , and that the tangent line of the curve at  $P_0$  is given by  $y = 0$ . This implies that  $c_{10} = c_{20} = c_{30} = c_{40} = 0$  in (1). Moreover, as  $2P_0$  is a divisor in the class  $\theta_0$  it follows that  $4P_0 = \text{div}(y) \frac{dx}{f_y}$ , the intersection divisor of the line  $y = 0$  with the curve, is a canonical divisor in the kernel of the Cartier operator. Given the above expression of the Hasse-Witt matrix this implies  $c_{12} = 0$ . With these normalizations the equation (1) for the curve simplifies to

$$f(x, y) = 1 + c_{01}y + c_{11}xy + c_{02}y^2 + c_{21}x^2y + c_{03}y^3 + c_{31}x^3y + c_{22}x^2y^2 + c_{13}xy^3 + c_{04}y^4.$$

The expression  $w_K$  is then given by

$$w_K = [(c_{21}^2c_{31} + c_{11}c_{31}^2)y]x^5 + [(c_{21}^2c_{22} + c_{31}^2c_{02})y^2 + (c_{21}^3 + c_{31}^2c_{01})y]x^4 + [(c_{11}c_{13}^2 + c_{03}^2c_{31})y^5 + (c_{01}^2c_{31} + c_{11}^3)y]x + (c_{13}^2c_{02} + c_{03}^2c_{22})y^6 + (c_{13}^2c_{01} + c_{03}^2c_{21})y^5 + (c_{01}^2c_{22} + c_{11}^2c_{02})y^2 + (c_{01}^2c_{21} + c_{11}^2c_{01})y.$$

If we successively subtract multiples of  $f(x, y)$  from the expression  $w_K$  so as to cancel in the resulting expressions the initial terms with respect to the lexicographic order with  $x > y$ , we obtain a remainder of the form

$$r = c_{31}^5x^3 + \dots.$$

This remainder must be a constant, and so  $c_{31} = 0$ , but this condition will result in a singularity at  $P_0$ , and so the result is proved.  $\square$

For the remainder we use other normalizations for such a curve. Now we bring the support  $P_0$  of the positive divisor in the canonical theta characteristic  $\theta_0$  to the origin  $(0 : 0 : 1)$  and force the tangent of the curve at  $P_0$  to be the line given by  $x = y$ . This tangent line intersects the curve at  $P_0$  with contact multiplicity 4, and thus never meets the curve again.

**Theorem 3.2.** *The isomorphism classes of curves of genus three having the canonical theta characteristic represented by a positive divisor supported at one point form an algebraic variety of dimension 4.*

*Proof.* In terms of the coefficients of the equation (1) defining the curve the normalizations stated above imply  $c_{00} = c_{10} + c_{01} = c_{20} + c_{11} + c_{02} = c_{30} + c_{21} + c_{12} + c_{03} = 0$ .

As  $2P_0$  is a divisor in the class  $\theta_0$  it follows that  $4P_0 = \text{div}(x + y) \frac{dx}{f_y}$ , the

intersection divisor of the line  $x = y$  with the curve, is a canonical divisor in the kernel of the Cartier operator. Given the above expression of the Hasse-Witt matrix this implies  $c_{21} + c_{30} = c_{03} + c_{12} = 0$ .

From the preceding proposition we can use a projective plane transformation in order to take one Weierstrass point to the location  $Q_1 = (1 : 0 : 0)$  and another to  $Q_2 = (0 : 1 : 0)$ . The projective automorphisms that fix these normalizations of the origin  $P_0 = (0 : 0 : 1)$ , of the tangent at the origin  $y = x$  and of the two infinite Weierstrass points  $Q_1 = (1 : 0 : 0)$  and  $Q_2 = (0 : 1 : 0)$  form a subgroup  $G$  of  $PGL_2(k)$  consisting of (classes of) matrices

$$G := \left\{ \begin{pmatrix} s_{00} & 0 & 0 \\ 0 & s_{00} & 0 \\ 0 & 0 & s_{22} \end{pmatrix} ; s_{00}, s_{22} \neq 0 \right\} / k^*. \quad (2)$$

As a consequence of the choice of points  $Q_1$  e  $Q_2$  we have  $c_{04} = c_{40} = 0$ . The tangent lines at these points (after the necessary homogenizations and dehomogeneizations) are, respectively,

$$c_{30}z + c_{31}y = 0 \quad \text{and} \quad c_{03}z + c_{13}x = 0.$$

These are Weierstrass points, and so these equations divide the quadratic parts appearing in the local expression of  $f$ . These divisibility conditions yield

$$\begin{aligned} c_{31}^2 c_{20} + c_{30}^2 c_{22} + c_{30} c_{31} c_{21} &= 0 & \text{and} \\ c_{13}^2 c_{02} + c_{03}^2 c_{22} + c_{03} c_{13} c_{12} &= 0, \end{aligned}$$

and these may be rewritten as

$$c_{31}^2 c_{20} + c_{30}^2 c_{22} + c_{30}^2 c_{31} = 0 \quad \text{and} \quad (3)$$

$$c_{13}^2 c_{02} + c_{03}^2 c_{22} + c_{03}^2 c_{13} = 0. \quad (4)$$

The result is now just parameter counting, once we observe that the two conditions above are algebraically independent. The curve is given by the equation  $f = f_1 + f_2 + f_3 + f_4 = 0$ , where  $f_i$  is the homogeneous part of degree  $i$  so that, with the chosen normalizations,

$$\begin{aligned} f_1(x, y) &= x + y \\ f_2(x, y) &= c_{20}x^2 + (c_{20} + c_{02})xy + c_{02}y^2 \\ f_3(x, y) &= c_{30}x^3 + c_{30}x^2y + c_{03}xy^2 + c_{03}y^3 \\ f_4(x, y) &= c_{31}x^3y + c_{22}x^2y^2 + c_{13}xy^3, \end{aligned}$$

is isomorphic, through a projective plane transformation given by an element of the group  $G$  described above, to the curve given by

$$f_1(x, y) + \alpha^{-1} f_2(x, y) + \alpha^{-2} f_3(x, y) + \alpha^{-3} f_4(x, y) = 0,$$

for  $\alpha = \frac{s_{00}}{s_{22}}$ . In these equations the conditions (3) and (4) have not yet been taken into consideration.  $\square$

#### 4. Curves with two Weierstrass directions towards the support of the canonical theta characteristic

We can ask the question of when the curve has two Weierstrass points  $Q_1$  and  $Q_2$  with Weierstrass directions towards  $P_0$ :

$$Q_1 \xrightarrow{W} P_0 \quad \text{and} \quad Q_2 \xrightarrow{W} P_0.$$

We can certainly use a projective plane transformation to bring these Weierstrass points to the chosen infinite locations  $(1 : 0 : 0)$  and  $(0 : 1 : 0)$ . This implies  $c_{30} = c_{03} = 0$ , and from (3) and (4) we deduce

$$c_{31}^2 c_{20} = 0 \quad \text{and} \quad c_{13}^2 c_{02} = 0.$$

On the other hand, if  $c_{31} = 0$  or  $c_{13} = 0$  then the infinite points  $Q_1$  or  $Q_2$  are singular, respectively, and so both  $c_{31}$  and  $c_{13}$  are non-zero, so that if  $c_{30} = c_{03} = 0$  then also  $c_{20} = c_{02} = 0$ .

If  $c_{30} = c_{03} = 0$  and  $c_{20} = c_{02} = 0$  then the equation of the curve simplifies to

$$C_{a,b,c} : \quad x + y + ax^3y + bx^2y^2 + cxy^3 = 0. \tag{5}$$

This family of curves were used in [RV] for displayed examples of minimal curves and eventually minimal curves.

**Remark 4.1.** *We left the details to the reader to check that a curve  $C_{a,b,c}$  as above is irreducible if and only if  $a + b + c \neq 0$ .*

The proof of the following result is straightforward:

**Proposition 4.2.** *The curve  $C_{a,b,c}$  is singular if and only if  $abc = 0$ .*

The intersection divisor with the infinite line is given by

$$Q_1 + Q_2 + Q_\delta + Q_{\delta+1},$$

where  $Q_\delta = (\frac{b}{a}\delta : 1 : 0)$  and  $Q_{\delta+1} := (\frac{b}{a}\delta + \frac{b}{a} : 1 : 0)$ , for  $\delta$  is any root of the Artin-Schreier equation  $t^2 + t + \frac{ca}{b^2} = (t + \delta)(t + \delta + 1) = 0$ . Under the hypothesis  $a, b, c \neq 0$  these points are all distinct. Moreover, the next result shows that these 4 infinite points are Weierstrass points with order sequence 0, 1, 3 and have Weierstrass directions towards the origin and there is no finite Weierstrass point with Weierstrass direction towards the origin.

**Theorem 4.3.** *In the family of curves  $C_{a,b,c}$  the Hasse-Witt invariant  $\sigma$  is 0 or 2 according to  $a = c$  or not. The rank of the Cartier operator is always 2.*

*In the curve  $C_{a,b,c}$  the origin is a Weierstrass point having order sequence 0, 1, 4 and its Weierstrass weight is 5, 8 or 20 according to  $a \neq c, a = c \neq b$  or  $a = b = c$ , respectively, and in any case it is the unique point with this Weierstrass weight. All 4 infinite points are Weierstrass points with order sequence 0, 1, 3, Weierstrass weight 1, having Weierstrass directions towards the origin and being the only ones with this latter property. There are other Weierstrass points with order sequence 0, 1, 4 only in curves  $C_{a,1,1/a}$  with  $a \neq 1$ , and these points have Weierstrass weight 4.*

*Proof.* The Hasse-Witt matrix of the curve  $C_{a,b,c}$  is given by

$$H = \begin{pmatrix} 0 & 1 & 1 \\ a & 0 & 0 \\ c & 0 & 0 \end{pmatrix}$$

and thus the Hasse-Witt invariant  $\sigma$ , which is the rank of the matrix

$$\begin{pmatrix} 0 & 1 & 1 \\ a & 0 & 0 \\ c & 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 & 1 \\ a^2 & 0 & 0 \\ c^2 & 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 & 1 \\ a^4 & 0 & 0 \\ c^4 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & a^2 + c^2 & a^2 + c^2 \\ a(a^4 + c^4) & 0 & 0 \\ c(a^4 + c^4) & 0 & 0 \end{pmatrix}$$

is given by

$$\sigma = \begin{cases} 2 & \text{if } a \neq c \\ 0 & \text{if } a = c \end{cases}$$

If  $a = c$  then the origin  $P$  is the unique point having 0, 1, 4 as order-sequence. In fact, if  $Q$  is another point with this order-sequence then we have  $4P \sim 4Q$  where  $\sim$  means linearly equivalent to. Since there are no 2-torsion points (the Hasse invariant is zero) we get  $2P \sim 2Q$ , but the curve is not hyperelliptic.

The numerator of the Wronskian for the curves  $C_{a,b,c}$  is given by

$$\begin{aligned} w_K &= f_x f_y (ax^2 + cy^2) + f_y^2 (axy + by^2) + f_x^2 (bx^2 + cxy) \\ &= bx^2 + by^2 + (a + c)xy + (ax^2 + cy^2) + (ax^2 + cy^2)^2 (x + y). \end{aligned}$$

The following hold:

$$\begin{aligned} \operatorname{div}(x) &= 2Q_2 + P_0 - [Q_1 + Q_\delta + Q_{\delta+1}] \\ \operatorname{div}(y) &= 2Q_1 + P_0 - [Q_2 + Q_\delta + Q_{\delta+1}]. \end{aligned}$$

Because of ([SV1] Theorem 1.5) the origin is a Weierstrass point with



weight greater than 2 having  $x$  as a local parameter, and the following expansions hold

$$\begin{aligned} y &= x + (a + b + c)x^4 + (a + c)(a + b + c)x^7 + \\ &\quad + ((a + c)^2(a + b + c) + (b + c)(a + b + c)^2)x^{10} + \dots \\ w_K &= (a + c)(a + b + c)x^5 + (a + b + c)^2(b + c)x^8 + \dots \end{aligned}$$

If  $a = c$  then the order of  $w_K$  at the origin is greater than 5, and the expansions are

$$\begin{aligned} y &= x + bx^4 + (b + a)b^2x^{10} + ab^3x^{13} + ab^4(a + b)x^{19} + \dots \\ w_K &= b^2(b + a)x^8 + b^4(b^3 + a^3 + b^2a)x^{20} + \dots \end{aligned}$$

If also  $b + c = 0$  (so that  $a = b = c$ ) then the order of  $w_K$  at the origin is 20, and the origin is the unique finite Weierstrass point. Summarizing we have:

$$v_{P_0}(w_K) = v_{P_0}(\mathcal{R}_K) = \begin{cases} 5 & \text{if } a \neq c \\ 8 & \text{if } a = c \neq b \\ 20 & \text{if } a = b = c \end{cases}$$

At the points  $Q_1$  and  $Q_2$  the following holds:

$$v_{Q_2}(w_K) = v_{Q_1}(w_K) = -5 \quad \text{and} \quad v_{Q_2}(\mathcal{R}_K) = v_{Q_1}(\mathcal{R}_K) = 1,$$

and so these points are Weierstrass points with orders 0, 1, 3 and weight 1.

The other infinite points  $Q_\delta$  and  $Q_{\delta+1}$  are Weierstrass points with orders 0, 1, 3 and weight 1, since taking the local parameter  $t = 1/y$  the local expansion of  $x$  begins as

$$x = \frac{b}{a}\delta t^{-1} + \dots,$$

and thus

$$ax^2 + cy^2 = \left( \frac{b^2}{a}\delta^2 + c \right) t^{-2} + \dots;$$

as  $\delta^2 + \delta = \frac{ca}{b^2}$  this first coefficient  $\frac{b^2}{a}\delta^2 + c = \frac{b^2}{a}(\delta^2 + \frac{ca}{b^2}) = \frac{b^2}{a}\delta$  is nonzero. At  $Q_\delta$ , for instance, the tangent is given by  $x + \frac{b}{a}\delta y = 0$ , and the intersection divisor of this tangent with the curve is given by  $3Q_\delta + P_0$ , so that  $Q_\delta$  and  $Q_{\delta+1}$  have Weierstrass directions towards the origin.

A line in the pencil through the origin has the equation  $y = \alpha x$ , which taken into the equation for  $C_{a,b,c}$  gives

$$x(1 + \alpha + x^3(a\alpha + b\alpha^2 + c\alpha^3)) = 0.$$

If  $\alpha \neq 1$  this equation does not have a multiple root, showing that no finite point of the curve other than the origin has its tangent passing through the

origin. As a consequence, the infinite points are the only ones in the curve with Weierstrass direction towards the origin.

Second coordinates of other Weierstrass points are given by the Sylvester resultant  $R(f, w_K)$  between  $f$  and  $w_K$ :

$$\begin{aligned} R(f, w_K) = & \\ = & y^{20}(a^2(c^4ab^4 + c^4b^5 + c^5b^4)) + y^{17}(a^2(ac^4b^3 + b^5c^3 + b^4c^4 + a^2c^3b^3)) + \\ & + y^{14}(a^2(b^7 + c^3a^4 + a^2c^5 + a^2b^4c + a^3c^2b^2 + a^2c^2b^3 + b^6a + b^4c^3)) + \\ & + y^{11}(a^2(b^5c + a^3c^2b + b^4a^2 + b^5a + c^2a^4 + a^2c^4 + b^4c^2 + ba^2c^3)) + \\ & + y^8(a^2(b^3a^2 + a^4c + b^2a^3 + a^2c^3)) + y^5(a^2(a^4 + a^2c^2 + ba^3 + ba^2c)). \end{aligned}$$

The roots of  $R(f, w_K)$  are the second coordinates of finite Weierstrass points, but the counting of multiplicities needs some care:  $R(f, w_K)$  has a multiple root in a higher Weierstrass point (that is, one having a higher weight in the ramification divisor or, equivalently (cf. [SV1] Theorem 1.5), having orders 0, 1, 4), but a multiple root of  $R(f, w_K)$  would also happen if 2 distinct Weierstrass points had the same second coordinate. Note that the origin is counted with multiplicity 5 if  $a \neq c$ , 8 if  $a = c \neq b$  and 20 if  $a = b = c$ .

If the curve has Weierstrass points with orders 0, 1, 4 other than the origin then the polynomial  $R(f, w_K)$  has other multiple roots. This situation is given by the discriminant of  $\frac{R(f, w_K)}{y^5}$ . To simplify the computation of this discriminant we set  $b = 1$ , which is allowed because of the action of  $G$ ; this discriminant is then given by

$$\text{disc} \left( \frac{R(f, w_K)}{y^5} \right) = (a + c)^2(a + c + 1)^2(1 + ac)^{12}.$$

The factor  $a + c$  is expected: the origin in this case has multiplicity greater than 5. The second factor is  $a + c + 1 = a + b + c$ , which is never zero. If the third factor is zero then  $a = \frac{1}{c}$ , and then

$$a^3 \frac{R(f, w_K)}{y^5} = (a^2 + a + 1)(y^3 + a(a + 1))(y^{12} + a^4(a + 1)).$$

If  $a^2 + a + 1 = 0$  then  $a \in \mathbb{F}_4 \setminus \mathbb{F}_2$ , but then  $c = \frac{1}{a} = a + 1 = a^2$  and hence  $a^2 + a + 1 = a + b + c = 0$ , which is against our moduli hypothesis. If  $a \notin \mathbb{F}_4$  then the three distinct roots of

$$y^{12} = a^4(a + 1)$$

are second coordinates of Weierstrass points with order sequence 0, 1, 4, all of them having weight four, and the three distinct roots of

$$y^3 = a(a + 1)$$

are second coordinates of Weierstrass points with order sequence 0, 1, 3, all of them having weight 1, as follows from direct computations.  $\square$

**Theorem 4.4.** *The isomorphism classes of curves of genus 3 having the canonical theta characteristic represented by a positive divisor supported at one point having 2 Weierstrass directions towards it form an algebraic variety of dimension two.*

*Proof.* There are exactly 4 Weierstrass points having Weierstrass directions towards the origin, and we may choose 2 of them to the normalized positions  $(1 : 0 : 0)$  and  $(0 : 1 : 0)$  in 12 ways. Having chosen these 2 infinite Weierstrass points, the group that fixes these normalizations is the group  $G$  described in (2) above. This group fixes the point  $P_0 = (0 : 0 : 1)$  and each point of the infinite line  $Z = 0$ , acting by homothety on finite points:  $(a : b : 1) \mapsto (\alpha a : \alpha b : 1)$ , where  $\alpha := \frac{s_{00}}{s_{22}}$ . Thus the curve  $C_{a,b,c}$  is isomorphic to the curve  $C_{\lambda a, \lambda b, \lambda c}$  for  $\lambda \neq 0$ .  $\square$

**Theorem 4.5.** *In the curve  $C_{a,b,a}$  all Weierstrass points other than the origin are simple Weierstrass points. If  $b = a$  the origin is the unique finite Weierstrass point, otherwise the other 16 Weierstrass points are 4 by 4 collinear. These 4 lines containing them are lines in the pencil of lines through  $(1 : 1 : 0)$ , and so is the canonical bitangent.*

*Proof.* If  $a = c$  then there are no other bitangents, and thus all Weierstrass points except the origin have orders 0, 1, 3 and weight one. Besides the infinite points, these Weierstrass points are given by the zeros of

$$\begin{aligned} w_K(x, y) &= a^2(x + y)^5 + (a + b)(x + y)^2 \\ &= (x + y)^2[a^2(x + y)^3 + a + b] \end{aligned}$$

As  $x \neq y$  away from the origin, these Weierstrass points occur in pairs  $(x : y : 1)$  and  $(y : x : 1)$ , the above equation being symmetric in  $x$  and  $y$ , as expected: if  $a = c$  then  $x \leftrightarrow y$  is an automorphism of the curve. The above equation gives a relation among the coordinates of Weierstrass points

$$x + y = \left( \frac{a + b}{a^2} \right)^{1/3}. \quad (6)$$

If  $a = b = c$  only the origin is a finite Weierstrass point, as we have seen. If  $a = c \neq b$  then  $\frac{a+b}{a^2}$  never vanishes and has 3 distinct cubic roots, and thus the other 12 Weierstrass points are on the 3 lines given by (6), occurring in 3 sets of 4 collinear points. These 3 lines are concurrent at the infinite point  $(1 : 1 : 0)$ , which is also a point in the canonical bitangent  $x = y$ .  $\square$

If  $a \neq c$  there are 3 non-canonical bitangents, corresponding to  $\alpha_0 + \alpha_1 x + \alpha_2 y = 0$  where

$$\begin{pmatrix} \alpha_0 \\ \alpha_1 \\ \alpha_2 \end{pmatrix}$$

is a solution of

$$H \begin{pmatrix} \alpha_0 \\ \alpha_1 \\ \alpha_2 \end{pmatrix} = \begin{pmatrix} 0 & 1 & 1 \\ a & 0 & 0 \\ c & 0 & 0 \end{pmatrix} \begin{pmatrix} \alpha_0 \\ \alpha_1 \\ \alpha_2 \end{pmatrix} = \begin{pmatrix} \alpha_0^2 \\ \alpha_1^2 \\ \alpha_2^2 \end{pmatrix}$$

(cf. [SV2] Proposition 3.3). Therefore, for each one of the 3 distinct roots of  $\alpha^3 = a + c \neq 0$ , there is one non-canonical bitangent given by

$$\alpha^{1/2} + a^{1/2}x + c^{1/2}y = 0.$$

These non-canonical bitangents are concurrent at the infinite point  $(c^{1/2} : a^{1/2} : 0)$ .

## References

- [K] K. Komiya, *Algebraic curves with non classical types of gap sequences for genus three and four*, Horoshima Math. Journal 8 (1978) 371-400.
- [NR] E. Nart, C. Ritzenthaler, *Non-Hyperelliptic Curves of Genus Three over Finite Fields of Characteristic Two*, arXiv:math/0312366v2 [math.NT] 2 Feb. 2004.
- [RV] J. E. A. Rodriguez, P. Viana, *Eventually Minimal Curves*, Bull. Braz. Math. Soc., New Series 36(1) (2005) 1-20.
- [St] H. Stichtenoth, *Algebraic Function Fields and Codes*, Springer Universitext, Springer Verlag, (1993).
- [SV1] K.-O. Stöhr, J.F. Voloch, *Weierstrass points and curves over finite fields*, Proc. London Math. Soc. 52 (1986) 1-19.
- [SV2] K.-O. Stöhr, J.F. Voloch, *A formula for the Cartier operator on plane algebraic curves*, Journal für die Reine und Ang. Math. 377 (1986) 49-64.
- [TP] O. P. La Torre, P. Viana, *Curves of Genus Three in Characteristic Two*, Communications in Algebra 33 (2005) 4291-4302.
- [V] A. M. Vermeulen, *Weierstrass points of weight two on curves of genus three*, Doctor Thesis, Universiteit van Amsterdam (1983).